

FQL
584
K3L4

UC-NRLF



D 2 535 260

Smithsonian Contributions to Knowledge.

THE

COLEOPTERA

OF

KANSAS AND EASTERN NEW MEXICO.

BY

JOHN L. LE CONTE, M. D.


WASHINGTON CITY:

PUBLISHED BY THE SMITHSONIAN INSTITUTION.

DECEMBER, 1859.

NEW YORK: D. APPLETON & CO.

EXCHANGE



Zoology Dept.

BIOLOGY
LIBRARY
6





UNIV. OF
CALIFORNIA





Illustration
of the
ENTOMOLOGICAL PROVINCES
of
NORTH AMERICA.

Prepared for the
Smithsonian Institution,
by
John L. Leconte, M.D.

SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE.

THE

COLEOPTERA

OF

KANSAS AND EASTERN NEW MEXICO.

BY

JOHN L. LE CONTE, M. D.

[ACCEPTED FOR PUBLICATION, OCTOBER, 1859.]



NOV 10 1859

f 9L584

KEL4

BIOLOGY
LIBRARY
6

COMMISSION

TO WHICH THIS MEMOIR HAS BEEN REFERRED.

C. ZIMMERMANN,
F. E. MELSHEIMER.

JOSEPH HENRY,
Secretary S. I.

PHILADELPHIA

COLLINS, PRINTER,
PHILADELPHIA.

INTRODUCTION.

THE present memoir is intended to give a catalogue of the Coleoptera thus far known to inhabit the middle eastern portion of the great central region of temperate North America. The boundaries of the province here treated of are as follows: north by the Missouri river, east by the meridian of the mouth of the Kansas or Platte river, south by about latitude 34°, and west by the main mass of the Rocky mountains.

It thus includes Kansas, a portion of Nebraska, and the eastern part of New Mexico. The eastern limit of this province is well defined; the other boundaries are indefinite, since it there fades imperceptibly into other provinces of the same great zoological district.

Before proceeding to consider the special material used in the preparation of this memoir, it will be proper to give a short sketch of the general results thus far obtained regarding the geographical distribution of Coleopterous insects in the territory of our republic.

The whole region of the United States is divided by meridional, or nearly meridional lines into three, or perhaps four, great zoological districts, distinguished each by numerous peculiar genera and species, which, with but few exceptions, do not extend into the contiguous districts. The eastern one of these extends from the Atlantic Ocean to the arid prairies on the west of Iowa, Missouri, and Arkansas, thus embracing (for convenience merely) a narrow strip near the sea-coast of Texas. This narrow strip, however, belongs more properly to the eastern province of the tropical zoological district of Mexico.

The central district extends from the western limit of the eastern district, perhaps to the mass of the Sierra Nevada of California, including Kansas, Nebraska, Utah, New Mexico, Arizona, and Texas. Except Arizona, the entomological fauna of the portion of this district west of the Rocky mountains, and in fact that of the mountain region proper, is *entirely* unknown; and it is very probable that the region does in reality constitute two districts bounded by the Rocky mountains, and southern continuation thereof.

The western district is the maritime slope of the continent to the Pacific, and thus includes California, Oregon, and Washington territories.

These great districts are divided into a number of provinces, of unequal size, and which are limited by changes in climate, and therefore sometimes distinctly, sometimes vaguely defined.

The Atlantic district may be divided into: 1, a northern province, including Maine, Eastern Canada, Nova Scotia, Newfoundland, etc., and extending west-

wardly from Lake Superior to Lake Winnipeg and Western Canada, which fades insensibly into the great Arctic district; 2, a middle province, limited westwardly by the Appalachian chain, and extending to Southern Virginia; 3, a western province, including Minnesota and the States of the valley of the Mississippi, as far as the State of that name; 4, a southern province, including the States south of Virginia and Kentucky; 5, a subtropical province, including the point of the peninsula of Florida; 6, a subtropical province, including the sea-coast of Texas.

The Central district, as far as known, may be thus divided: 1, a northern province, comprising the regions north of the Missouri, the plains of the Saskatchewan, etc.; 2, a middle eastern province, divided into two subprovinces, including: *a*, Kansas, and Nebraska; *b*, northeastern New Mexico; 3, a southeastern province, including Texas, with the exception of province six of the Atlantic district; 4, a southwestern province, including the upper part of the valley of the Gila; and 5, a south-southwestern province, including the lower Gila and Colorado. The unexplored portions of this district will indicate middle western, and northwestern provinces, or perhaps the necessity of constituting with them and the southwestern province a district to be called the Interior district.

The Pacific district may be divided as follows: 1, a hyperborean province, consisting of Sitka and the neighborhood; 2, a northern province, including Eastern Oregon and Washington Territories; 3, a middle province, including California, probably as far south as Santa Barbara; 4, a southern province, including California from Santa Barbara to San Diego, extending to the crest of the Sierra. Southern, or lower California is also, perhaps only in part, a province of this district;¹ but, as yet, no collections of magnitude have been received therefrom. Other provinces will, from the peculiar method of distribution of species in that portion of America, be defined when more full collections are made, but at present cannot be indicated.

At the north, the Atlantic and Central districts seem to merge imperceptibly together, about the valley of the Athabasca, and Winnepeg rivers, and finally to disappear in the limited Arctic fauna; the hyperborean province of the Pacific district also fades into this Arctic fauna, without, however, losing itself so perfectly in the northern provinces of the other districts. We have thus evidence that the American Arctic district may be divided into two provinces, an eastern and a western.

At the south, the Atlantic district merges through Florida into the Caribbean tropical province, and through maritime Texas into the Mexican lower eastern province. In the same direction the Central district merges into the Mexican upper or central province, and the Interior district, towards the Gulf of California, into the Mexican western province. Regarding the southern affiliations of the Pacific district we know absolutely nothing; scarcely a single species found at San Diego has been found in Mexico.

¹ A few species, collected by John Xantus, Esq., at Cape San Lucas, though all new, indicate a greater resemblance to the fauna of the lower Colorado, than to that of maritime California; this province may therefore be found eventually to belong to the interior district.

The method of distribution of species in the Atlantic and Pacific districts, as already observed by me in various memoirs, is entirely different. In the Atlantic district, a large number of species are distributed over a large extent of country; many species are of rare occurrence, and in passing over a distance of several hundred miles, but small variation will be found in the species obtained. In the Pacific district, a small number of species are confined to a small region of country; most species occur in considerable numbers, and in travelling even one hundred miles, it is found that the most abundant species are replaced by others, in many instances very similar to them; these small centres of distribution can be limited only after careful collections have been made at a great number of localities, and it is to be hoped that this very interesting and important subject of investigation may soon receive proper attention from the lovers of science on our Pacific shores.

In the Central district, consisting, as it does to a very large extent, of deserts, the distribution seems to be of a moderate number of species over a large extent of country, with a considerable admixture of local species; such at least seems to be the result of observations in Kansas, Upper Texas, and Arizona.

For the purpose of enabling these investigations to be carried on in future with less labor, I have caused the catalogues of the present memoir to be printed separately; the small size of the catalogue of species from Eastern New Mexico, will call attention to the necessity of procuring more material from that region; while the asterisks affixed to the species which have occurred in both sub-provinces, will give the results thus far obtained in geographical distribution.*

For many years in the early history of entomology in the United States, the Coleoptera of Kansas were as well known, and even more fully described than those found in the Atlantic States. They form, indeed, the subject of one of the earliest and most extensive of the valuable contributions to entomology made by Say.

Having, in the year 1845, made a journey along the Platte river to Fort Laramie, thence near the base of the mountains to the Arkansas, returning by that river and the Santa Fé road, I was enabled to follow nearly in the footsteps of Say, and had the singular good fortune to recover nearly all the species described by him, and of which the types had been destroyed to such an extent that scarcely an authentic specimen remained.

Of the species described by him, and not obtained by me from this region, I have introduced those which remain unknown to me, marking them thus (*fide* Say); the others I have excluded, as I possess them from Missouri, the locality in most

* The student will also consult in reference to the Coleopterous fauna of the Central and Pacific districts: 1. My report on the insects collected along parallel 47°, in Pacific R. R. Explorations and Surveys, vol. xi., which includes a list of the species found on the Pacific Slope, north of San Francisco; an appendix to the same in Proceedings of the Academy of Natural Sciences of Philadelphia, October, 1859.

2. Catalogue of Coleoptera of the region adjacent to the boundary line between the United States and Mexico; Journal of the Academy of Natural Sciences of Philadelphia, second series, vol. iv., No. 1.

3. Catalogue of the Coleoptera of Fort Tejon, California; Proceedings of the Academy of Natural Sciences of Philadelphia, February, 1859.

instances given by Say, and have no evidence that they are found within the region herein considered.

The results of my own labors have been measurably increased by the following collections, which I owe to the kindness of friends, and to the Smithsonian Institution:—

1. Collections brought by Lieut. Beckwith's expedition; among which were two new species; *Cleonus angularis*, and *Cœlocnemis punctatus* Lec. (Proc. Acad. Nat. Sc. Philad., VII, 225). The genus *Cœlocnemis* was not previously known east of California, where it is represented by several species; suspecting that it was collected inside of the Great Basin, I have excluded it from the catalogues here given.

2. A large number of specimens collected in Eastern Kansas, by Mr. M. Burke, and presented to me through Dr. John Torrey.

3. A very large collection made at Fort Riley, Kansas, by Dr. W. A. Hammond, U. S. A., and John Xantus, Esq.

4. Collections from the mouth of the Yellowstone river, and from the Loup Fork of the Platte, made by Dr. F. V. Hayden, and received from the Smithsonian Institution.

5. A collection made by Dr. Wm. A. Hammond, on the route from Fort Riley to Bridger's pass.

6. Similar collections made by Dr. John G. Cooper, received from the Smithsonian Institution.

The materials received from Eastern New Mexico are much more scanty, and are as follows:—

1. Four or five hundred specimens collected near Santa Fé, by Mr. Fendler, and procured for me by the kind offices of Dr. Engelmann.

2. Specimens collected by Dr. Wislizenus, on a journey from Santa Fé to Chihuahua; though few in number, these specimens were very interesting in character, and I have availed myself of the present opportunity to figure some of them, though they are possibly not found within this province of the Central district, but belong rather to the southwestern province.

3. A small, but very interesting collection made in the vicinity of Santa Fé, by the late R. C. Kern, and given me by Prof. S. S. Haldeman.

These materials were used in preparing articles for the reports of the expeditions of Capt. E. G. Beckwith, and Capt. J. Pope, U. S. A., in Pacific R. R. series, but were excluded from want of room, and are now incorporated together, with the addition of several new species since obtained from various sources.

The six provinces of the Atlantic district are marked on the map by red tints; the two provinces and two subprovinces of the Central district proper by green; the two southern provinces of the supposed Interior district by brown; and the four provinces thus far ascertained of the Pacific district by blue tints.

COLEOPTERA OF KANSAS AND EASTERN NEW MEXICO.

AMBLYCHILA SAY.

A. cylindriformis SAY, Trans. Am. Phil. Soc. IV, 409.

The specimen figured (Tab. II, fig. 1), was found by Capt. Pope on the Llano Estacado; it is very large (1.28 inches long), and in form is altogether similar to the specimen figured by me in the Proceedings of the Academy of Natural Sciences, vol. vii. p. 32. The punctures of the elytra are however much stronger, being in fact hardly smaller than those which are scattered in irregular series. The costæ of the elytra are more elevated, and the internal one extends fully to the posterior declivity. The legs are thicker, and the posterior tibiæ are entirely straight. Two other specimens were collected by Dr. Wm. A. Hammond, U. S. A., on the Platte river, about 100 miles above Fort Kearny. They agree in sculpture with the specimen from the Llano Estacado; one of them, which I consider as a male, has rather narrower elytra than the other, and has also the thorax more rounded on the sides: otherwise there is no difference between them. A larger series of specimens will indicate the nature of the strongly-marked differences of sculpture.

As confirming the correctness of the view expressed by Lacordaire regarding the identity of *A. cylindriformis* with the Californian *A. Piccolomini*, I may add, that having sent to Baron Chaudoir a figure and description of the labrum of the specimen found at Fort Union by Major Sibley, I was informed by that distinguished entomologist that he can find no appreciable difference between the figure and the labrum of the Californian specimen now in his cabinet. The figure given by M. Reiche is therefore incorrect, as already surmised.

MICRIXYS Lec.

M. distinctus, rubens, flavo-pilosus, capite thoraeque grosse punctatis, hoc transverso obovato, postice valde, angustato, lateribus valde rotundatis, elytris dorso subdepressis, striis valde cribratis, macula transversa laterali ad medium ornatis, apice summo, pedibus antennisque piceis, his articulo 1mo rufo. Long. .3. Tab. II, fig. 2.

Micrixys distinctus Lec. Proc. Acad. Nat. Sc. VII, 220.

Panagaeus distinctus HALD. Stansbury's Expedition to Great Salt Lake, 373.

Eugnathus || *distinctus* Lec. Trans. Am. Phil. Soc. X, 375.

New Mexico: one specimen found by the late Richard Kern, Esq. This genus, unfortunately established under a preoccupied name, differs remarkably from *Panagaeus* by the head not being constricted into a neck posteriorly, and by the mandibles being thick, dilated, obtuse, and deflexed at the apex.

CYMINDIS LATR.

C. cribrata, nigro-picea, setis brunneis pilosella, capite thoraceque cribratim punctatis, hoc tenuiter marginato convexo, latitudine longiore, postice modice angustato et lateribus parum sinuato, angulis posticis obtusissimis, elytris cyaneo-micantibus, striis foveatim punctatis, interstitiis uniseriatim punctatis, antennis palpis pedibusque ferrugineis. Long. .36.

Nebraska: one specimen collected by Mr. Burke. Resembles in character *C. pilosa*, but the thorax is longer, and less rounded on the sides anteriorly: the punctures of the striæ of the elytra are less closely placed, and those of the interstices almost form single rows. *C. neglecta* Hald. is smaller, and has the head almost smooth.

ANISODACTYLUS DEJ.

A. chalcus, oblongus, elongatus, supra æneus, thorace latitudine vix brevior, punctulato, parce punctato, lateribus postice vix sinuatis, angulis posticis rectis, basi utrinque punctato, profundius impresso, elytris striatis, interstitiis planis alterutim parce punctatis; subtus nigro-æneus, antennæ basi rufis; tibiis anticis calcare extrorsum paulo dilatato. Long. .37.

Santa Fé, Mr. Fendler. Allied to *A. alternans* Lec. (Ann. Lyc. Nat. Hist. 5, 184), but differs in color, and in the less dense punctuation of the alternate spaces between the striæ of the elytra.

HARPALUS LATR.

H. impotens, piceo-niger, oblongus, thorace transverso, lateribus rotundatis, margine angusto reflexo, angulis posticis rotundatis, basi utrinque vage foveato, versus angulos subdepresso, elytris ad marginem subtiliter pubescentibus, striis 2nda, 5ta, 7maque parce punctatis, interstitiis fere planis, epipleuris, ano pedibusque piceis, antennæ pedibusque rufo-piceis. Long. .38.

LEC. Journ. Acad. Nat. Sc. Philad. 2d ser. IV, 14.

One specimen found at El Paso by Dr. Thos. H. Webb. This species is narrower than *Harpalus (Selenophorus) stigmus*, and *iripennis*, and is readily distinguished by the rounded angles of the thorax and by the elytral punctures being smaller than in the first mentioned species.

H. oblitus, oblongus, piceo-niger, nitidus, thorace latitudine sesqui brevior lateribus rotundatis, postice subangustato, angulis posticis obtusis, margine piceo postice anguste subexplanato, basi utrinque vage impresso punctulato et rugoso, elytris striis tenuibus, secunda unipunctata; subtus, antennæ palpis pedibusque ferrugineis. Long. .5.

One male, Santa Fé, Mr. Fendler. Resembles *H. compar* Lec. (*pennsylvanicus* † Say; *bicolor* † Dej.) in general appearance, but by the thorax being slightly narrowed at base, with only a few small punctures around the basal impressions, it is abundantly distinct. Fully matured specimens will probably be found to have the under surface of the body dark colored.

H. fallax, oblongus, piceo-niger, nitidus, thorace latitudine sesqui brevior lateribus rotundatis, ante medium subangustato, angulis posticis rectis subrotundatis, margine laterali postice subexplanato, cum basi tota punctulato, ad basin utrinque late foveato, elytris striis bene impressis, 2da unipunctata; antennæ, palpis, pedibusque ferrugineis. Long. .42.

Santa Fé: Messrs. Kern and Fendler. The elytra are slightly sinuate at tip, and those of the female are dull. This species is of the size of *H. herbivagus* Say, but the thorax is distinctly narrowed before the middle, and is much more punc-

tured at base. A very similar nondescript species from New Jersey was kindly given me by Mr. Guex; it differs chiefly by the thorax being broader, with the sides less rounded and less narrowed anteriorly. I have named it *H. viluus*.

H. desertus, oblongus, piecus nitidas, thorace latitudine brevior, lateribus rotundatis vix explanatis, antice posticeque subangustato, angulis posticis subobtusis, subrotundatis, ad basin utrinque punctulato et foveato, elytris ad apicem obtuse rotundatis, striis bene impressis, 2nda unipunctata; antennis palpis pedibusque ferrugineis. Long. '3.

One immature female: Mr. Fendler. Quite different from any other species known to me, although having somewhat the form of *H. megalcephalus* Lec.

H. stupidus, oblongus subovalis, convexus, ater, thorace latitudine plus sesqui brevior, lateribus tenuiter marginatis antice rotundatis, postice subrectis fere parallelis, angulis posticis rectis vix rotundatis, fovea basali utrinque haud profunda, basi tota punctulata, elytris (feminae subopacis) striis impunctatis, interstitiis vix convexis, apice sinuatis, antennis pedibusque obscure ferrugineis. Long. '46.

One specimen from route to Fort Bridger: Mr. Drexler. More convex than *H. erythropus*, with the sides of the thorax not flattened, and the posterior angles less obtuse. From these differences in form, this species has somewhat the appearance of *H. funestus* Lec., which however has black feet, and is otherwise quite different.

DICÆLUS BON.

D. laevipennis, oblongo-ovalis, violaceo-niger, thorace antrorsum valde angustato, lateribus rotundatis, elytris tenuiter punctato-striatis, carina humerali ante medium desinente. Long. '7. Tab. I, fig. 1.

LEC. Annals of the Lyceum of Natural History of New York, IV, 421.

Platte River Valley, near Chinney Rock. Very different from all other species yet known, by the striæ of the elytra being indicated only by rows of punctures, which are less obvious towards the apex and sides.

NOMARETUS LEC.

N. cavicollis, æneo-purpureus, nitidus, thorace obovato, postice valde angustato, sulco dorsali profundo, antice disperse punctato, postice latius concavo, profunde impresso et punctato, elytris striis utrinque 12 fortiter arenatis. Long. '45.

One specimen, Fort Riley: Mr. John Xantus. Resembles *N. fissicollis* and *N. bilobus* (*Cychrus bilobus* Say), but differs from both by the anterior half of the thorax being covered with scattered punctures, and the posterior half being broadly concave; as in those species the posterior angles are obtuse.

CALOSOMA FABR.

C. luxatum, nigrum subnitidum, capite subtilius punctato-rugoso, thorace capite sesqui latiore, latitudine fere triplo brevior, lateribus valde rotundatis, postice angustiore confertim punctato-rugoso, basi late rotundatim emarginato, utrinque vage impresso, elytris rotundato-ovatis, thorace latioribus, striis tenuibus rugisque transversis imbricatim insculptis, versus apicem confuse rugosis. Long. '64—'68. Tab. I, fig. 2.

SAY, Journ. Acad. Nat. Sc. of Phila. III, 149; (nec Dej. Sp. Gen. II, 196).

Callisthenes luxatus LEC. Annals of the Lyceum of Nat. Hist. V, 200.

Carabus luxatus LEC. Ann. Lye. Nat. Hist. IV, 445.

Platte river: on one occasion seen running on the ground in large numbers. Wider and more robust than the other North American species.

C. striatulum, nigrum, capite thoraceque nitidis leviter rugosis, hoc brevi, versus latera et basin rugoso et punctato, postice paulo angustato, ad basin late emarginato, elytris thorace latioribus, subrotundatis, dense striatis, interstitiis imbricatis, versus apicem tuberculis parvis confertis exasperatis. Long. .65.

Milk river: one specimen found by Dr. Suckley, U. S. A., attached to the North Pacific R. R. Expedition under Gov. J. J. Stevens; Utah, Mr. Drexler. This species closely resembles *C. Zimmermanni* Lec., which is properly an Oregon species, but differs in the head and thorax being much less punctured; also in the thorax being less narrowed towards the base, and in the striæ of the elytra being more distinct.

Body black; head shining, finely but sparsely wrinkled and punctured, with the frontal impressions moderately deep. Thorax shining, about one-half wider than the head, twice as wide as its length, rounded on the sides anteriorly, obliquely but only moderately narrowed to the base, which is broadly emarginate, with prolonged basal angles; the middle of the base is, however, truncate, and not concave as in *C. luxatum*; the sides are narrowly margined, the dorsal line is distinct; the sides are densely rugosely punctured, but the sculpture becomes indistinct on the disc. The elytra are one-half wider than the thorax, about one-fourth longer than their width, covered with close set striæ, the outer ones of which are indistinct; the interstices are transversely rugose, the rugæ being deeper at the sides and apex, so that the surface is there covered with small rounded tubercles.

ILYBIUS ER.

I. Laramacus, elongatus ovalis, antice paulo obtusus, nigro-æneus, undique subtiliter dense reticulatus, fere opæus, elytris striola submarginali, maculaque majuscula anteaicali pallidis, ore pedibusque anterioribus piceo-rufis, pedibus posticis nigro-piceis. Long. .37—41.

Fort Laramie: narrower and less convex than our species found in the Atlantic States, with the sides of the thorax less rapidly converging in front: resembling more nearly in form the *I. 4-maculatus* of Oregon; it is, however, less oval than that species, with the sides less rounded, and the anteaical spot larger. The specific differences in this genus are very unsatisfactory, and depend entirely upon slight modifications in form, which cannot be clearly expressed in a description.

AGABUS LEACH.

A. clavatus, elongato-ovalis, modice convexus, piceus nitidus, obsolete punctulatus, capite, thoracis elytrorumque lateribus sensim rufo-piceis, pedibus antennisque ferrugineis, his articulis externis dilatatis, 9-11 nigro-piceis. Long. .34.

Three males from Loup Fork of the Platte: Dr. Hayden. A most interesting addition to our fauna, being the analogue of *A. serricornis* of Europe. The 5th, 6th, and 7th joints of the antennæ are gradually wider; the 8th, 9th, and 10th are subrectangular, wider than long, and as wide as the 7th; the 11th is oval and pointed, narrower than the 10th.

A. griseipennis, elongato-ovalis, parum convexus, æneo-niger, supra dense (femine fortiter) reticulatus, elytris luteo-griseis base margineque pallidioribus, antennis palpisque flavis, pedibus ferrugineis, femoribus nigro-maculatis; thorace eum elytris angulum haud formante. Long. .36.

Fort Laramie, one female. Differs from the species of somewhat similar color inhabiting the United States by its more elongate form, gradually tapering both before and behind; the thorax continues accurately the outline of the elytra. The reticulations are very dense, and somewhat longitudinal, especially at the base of the elytra. The Californian *A. lutosus* Lec. approaches it very closely, and it may in fact prove to be the other sex of that species, of which I possess at present only males. The form is, however, somewhat narrower.

A. obliteratus, ovalis, elongatus parum convexus, æneo-niger, nitidus supra subtilissime, vix conspicue reticulatus, elytris luteo-piceis, marginibus pallidioribus, seriebus punctorum solitis valde distinctis, ad basin thorace paulo latioribus, antennis palpis tibiis tarsisque anterioribus ferrugineis. Long. .32.

Fort Laramie, three specimens. The sculpture of the female is hardly more conspicuous than that of the male. It is sufficiently distinguished from our other species by the characters above given.

A. spilotus, ovalis, æneo-niger nitidus, parum convexus, vix subtilissime reticulatus, thorace eum elytris angulum haud formante, elytris seriebus solitis distinctis, lineola utrinque submarginali, guttaque subapicali pallidis, antennis palpisque rufis, tarsi anterioribus rufo-piceis. Long. .35.

Two females, Fort Laramie. Resembles *A. obtusatus*, but is less dilated, and differs very much by the reticulation being so minute as to be scarcely visible even with a high magnifier. The spots are not very obvious.

ANISOMERA BRULLÉ.

A. cordata, supra æneo-picea, elongato-ovalis depressa, subtilissime reticulata, thorace brevi, postice angustato et lateribus sinuato, elytris thorace vix latioribus, postice non dilatatis; subtus nigra, ore antennis pedibusque rubro-piceis. Long. .45. Tab. II, fig. 3.

LEC. Proc. Acad. Nat. Sc. VI, 226.

Santa Fé; Mr. Fendler. Varies in having black feet; the sides of the thorax are strongly rounded in front, and subsinuate behind; the base is as wide as the apex, the anterior dilatation being produced by the curvature of the sides; the anterior angles are acute, the posterior ones rectangular. The elytra are very little wider than the thorax, regularly elongate elliptical, with the usual series of punctures becoming irregular towards the tip.

By comparing the figure here given with the South American *A. bistriata* Brullé (figured in Dejean's Icon. Col. Europe, V, tab. 23, fig. 5), it will be seen that they correspond very closely. The middle lobe of the mentum of our species is broad, short, and very obsoletely sinuate; in the generic description given by Aubé, the mentum of *A. bistriata* is said to have the middle lobe slightly prominent in the middle. This character needs confirmation, as it is hardly possible that two species so closely related should differ in this particular. Though distinct in form, this genus is separated from *Agabus* by very trifling differences.

SILPHA LINN.

S. bituberosa, ovalis depressa, nigra opaca, dense punctata, thorace latitudine plus duplo brevior, lateribus anguste marginatis valde rotundatis late planatis, postice utriusque late oblique impresso, elytris fortiter marginatis, apice conjunctim rotundatis, costis tribus postice paulo abbreviatis, internis duabus parum distinctis, externa tuberculum posticum transeunte. Long. .5.

One specimen found near Fort Bridger, Mr. Drexler; belongs to the division of the genus having the 8th—11th joints of the antennæ about one-half wider than those which precede; the 8th to 10th are not much wider than long; the 2d is quite as long as the 3d.

S. truncata, oblonga, atra subopaca, capite thoraceque dense aciculato-punctatis, hoc antrorsum valde angustato, ad basin trisinuato, postice utriusque versus latera oblique impresso, elytris lateribus fortius marginatis, subtilius punctatis, ad dodrantem callo discoideo parum distincto utriusque notatis, ad apicem late truncatis; antennarum articulo 3io vix longiore. Long. .5—'67. Tab. I, fig. 3.

SAY, Journ. Acad. Nat. Sc. III, 193. LEC. Proc. Acad. Nat. Sc. VI, 278.

Platte river, near the Forks. A very distinct species; the specimen figured is a male: the abdomen of the female projects far beyond the elytra.

CARPOPHILUS LEACH.

C. apicalis, oblongus, piceo-niger, minus subtiliter punctatus et flavo-pubescent, thorace modice convexo, latitudine haud sesqui brevior, lateribus marginatis rotundatis, elytris thorace sesqui longioribus, rufis, circa scutellum et ad apicem extrorsum oblique infuscatis, pedibus antennisque testaceis. Long. .1.

Platte river: found also in Georgia; narrower and more convex than is usual in this genus.

C. carbonatus, oblongus subdepressus, piceo-niger, parcius pubescens et punctatus, thorace latitudine plus sesqui brevior angulis omnibus rotundatis, ad apicem vix emarginato, lateribus late rotundatis tenuiter marginatis, elytris alutaceis obsolete punctulatis, thorace duplo longioribus, antennis basi piceis. Long. .1.

Found on the Platte river and at Lake Superior.

MELIGETHES KIRBY.

M. ruficornis, oblongus subconvexus, nigro-viridis, dense subtiliter punctatus, breviter cinereo-pubescent, thorace antrorsum angustato, lateribus rotundatis anguste marginatis, pone medium subfoveatis, angulis posticis subrectis, antennis tarsis tibiisque rufis, his anticis subtiliter serratis, posterioribus latioribus ultra medium et ad apicem ciliatis; unguibus simplicibus, prosterno postice anguste rotundato. Long. .11.

Platte river, not rare. Differs by obvious characters from all European species thus far described, and were it not that the prosternum is rounded posteriorly, it would enter Erichson's division B, (Ins. Deutschl. 173.)

M. sævus, oblongus subconvexus, niger, dense subtiliter punctatus, breviter cinereo-pubescent, thorace antrorsum angustato, lateribus rotundatis anguste marginatis, angulis posticis rotundatis, margine basali undulato, antennis basi piceis; tibiis anticis 5-dentatis ad basin serratis, intermediis extrorsum emarginatis et posticis dense spinulosis; unguibus simplicibus, prosterno postice late rotundato. Long. .11.

One specimen; Platte river; belongs to Erichson's division C, (loc. cit. 179.)

HISTER LINN.

II. instratus, quadrato-ovalis, niger nitidus, thorace bistriato, lateribus dense flavo-ciliatis, elytris stria suturali utrinque abbreviata, tribus externis integris, marginali brevissima, rudimentalique ad basin interna notatis, macula utrinque lunata maxima rubra ornatis; tibiis anticis bidentatis, posticis compressis biserialim spinulosis, femoribus posticis majoribus rubro-tinctis. Long. ·24—·32.

Platte river: Intermediate between *H. arcuatus* Say, and *H. biplagiatus* Lec., having the elytral spot and compressed posterior tibiæ as in the latter, while by the very densely ciliate margin of the thorax, the thick and sometimes reddish posterior thighs, and the more numerous spines of the posterior tibiæ it is evidently related to the former. The outer series of spines is composed of numerous spines placed irregularly in nearly a double row, while in *H. biplagiatus*, they are distant and regular; the posterior tibiæ are slightly less compressed than in that species.

II. nubilus, rotundato-ovalis, niger, nitidus, thorace bistriato, stria externa margini approximata, ante medium abbreviata, elytris striis externis 4 integris, 5ta postica brevi, suturali ante medium antice abbreviata, epipleuris striis duabus punctatis, striolaque brevi inferna; mesosterno fere truncate; pygidio sat dense punctato, tibiis anticis sub-5-dentatis. Long. ·25.

One specimen: Platte river. This and the next are very distinct from any of our other species, in which the mesosternum is not distinctly emarginate, and the elytral marginal striæ wanting.

II. pollutus, rotundato-ovalis, niger nitidus, thorace bistriato, stria externa margine approximata ante basin abbreviata, elytris striis externis 3 integris, 4ta ad medium postice abbreviata, 5ta obsoleta, suturali utrinque parum abbreviata, epipleuris 3-striatis; mesosterno fere truncate; pygidio parcius punctato, tibiis anticis sub-4-dentatis. Long. ·20.

Kansas and New Mexico.

SAPRINUS LEACH.

S. spurcus, quadrato-rotundatus, æneo-niger, nitidus, capite punctulato, thorace lateribus dense punctatis, antice vix obsolete impresso, margine laterali sublævi, elytris striis dorsalibus æqualibus ad medium abbreviatis, interna cum suturali integra connexa, marginali brevi obliqua cum humerali fere juncta; punctatis postice subaciculatis, spatio sentellari quadrato, margineque laterali lævibus, epipleuris bistriatis punctatis, tibiis anticis sub-6-dentatis. Long. ·14.

One specimen: Platte river. Belongs to my division 4 (Proceedings Acad. Nat. Sc. Phila. VI, 40), having the head not margined in front, the epipleuræ bistriate, the prosternum transversely convex, with an anterior fovea on each side, and the prosternal striæ parallel, abbreviated in front. It is closely related to *S. lugens*, but is very considerably smaller, with the punctures of the elytra less aciculate, the thorax scarcely impressed, and with the punctures of the sides not extending to the lateral margin, which is only punctulate.

S. parumpunctatus, rotundatus, nigro-æneus nitidus, parce punctulatus, thorace lateribus et basi confertim punctato, antice haud impresso, elytris striis paulo pone medium abbreviatis, æqualibus, 4ta cum suturali integra conjuncta, marginali brevi, postice haud dense punctato, pedibus rufis, tibiis anticis denticulatis. Long. ·13.

One specimen: Platte river. Resembles *S. conformis*, and *placidus*, but differs, besides slighter characters above mentioned, by the punctures of the posterior

portion of the elytra being smaller and less deep. Belongs to my group 6, having the prosternum transversely convex, foveate each side in front, with the striæ remote and divergent; the head not margined in front, and the epipleuræ bistrate.

S. pratensis, rotundatus, nigro-æneus, vel nigro-cupreus, nitidus, thorace lateribus et basi confertim punctato, antice haud impresso, elytris striis ad medium abbreviatis æqualibus, 4ta cum suturali conjuncta, marginali brevi, dimidio postico dense punctatis, extrorsum fere aciculatis, punctis inter striam humeralem et 2ndam ad basin, extensis; pygidio dense punctato, pedibus rufis, tibiis anticis denticulatis. Long. 13—20.

Platte river, and Fort Riley, abundant. I was inclined to believe this to be *S. orbiculatus* Marseul found in Texas, but the punctures extending to the base are not entirely outside of the dorsal striæ, but occupy also the spaces as far as the 2d stria; these spaces are sometimes also slightly rugous. The prosternum is subcarinate, margined with the striæ divergent; it therefore belongs to my division 7, (Proc. Acad. VI, 40.)

PHILEURUS LATR.

P. valgus. (Tab. II, fig. 4.)

On comparing specimens from Georgia, Missouri, Texas, and New Mexico, forming *P. castaneus* Hald., with one from Brazil in the collection of the Academy of Natural Sciences, I can perceive no difference whatever. Burmeister has observed that the species extends its range from South America into Mexico, but adds that he has not seen specimens from the regions north of that country. Individuals occur of a dark brown color, but I suspect these to be immature: the usual color is a full black.

POLYPHYLLA HARRIS.

P. decemlineata, picea, dense luteo-squamosa, clypeo maris valde concavo, antice latiore recte truncato (femine lateribus parallelis apice sinuato), capitis lateribus albopilosis; thoracis canaliculati vittis tribus, scutello medio, elytrorum sutura vittis utrinque tribus humeralique brevi cum externa confluyente niveo-squamosis; subtus niveo-squamosa, pectoribus longe luteo-villosis, pedibus ferrugineis parcius squamosis (tibiis anticis maris bidentatis, femine tridentatis). Long. 1.0—1.4. Tab. I, fig. 6, (a, tib. ant. femine.)

LEC., Proc. Acad. Nat. Sc. VII, 218; Journ. Acad. Nat. Sc., 2d ser. III, 230.

Melolontha 10-lineata SAY, Journ. Acad. Nat. Sc. III.

Kansas, New Mexico, Texas, California, Oregon. The thorax has occasionally a round whitish spot near each side.

THYCE LEC.

T. squamicollis, ferrugineo-picea, capite thoraceque confertissime punctatis, pilis depressis squamiformibus pallidis obsitis, elytris nitidis punctatis et rugosis, brevissime parce pubescentibus, pygidio abdomineque squamulosis, pectore valde lanuginoso. Long. 1.03. Tab. II, fig. 5.

LEC. Journ. Acad. Nat. Sc. 2d ser. III, 232.

One female found at Albuquerque, by Dr. T. Charlton Henry, U. S. A. Reddish brown above and beneath; head densely punctured, front nearly straight anteriorly, with the margin broadly reflexed, covered with small pale yellowish scales. Thorax convex, narrowed in front, much rounded and subserate on the sides, broadly but feebly channelled in the middle, vaguely impressed each side near the anterior angles

which are acute; densely punctured, covered thinly with yellowish scales like those on the head; scutellum densely pubescent with a medial glabrous line. Elytra shining, not densely punctured, with rugæ as in most of the species of *Phyllophaga*; each puncture furnished with a pale short hair. Pygidium densely, abdomen sparsely covered with small pale scales. Pectus with dense long yellow hair. Anterior tibiae three-toothed; middle and posterior tibiae with two sharp teeth near the middle.

LACHNOSTERNA HOPE.

L. lanceolata, pieca (mas oblonga, femina crassa), supra squamulis lutescentibus dense tecta haud punctata, clypeo reflexo vix emarginato, thorace lateribus antice valde rotundatis, angulis posticis reectis, linea dorsali subglabra, elytris vittis utrinque tribus indistinctis subglabris; subtus dense sordide pubescens, antennis pedibusque obscure ferrugineis. Long. .53—.68. Tab. I, fig. 5(♀).

LEC. Journ. Acad. Nat. Sc. 2d ser. III, 237.

Melolontha lanceolata SAY, Journ. Acad. Nat. Sci. III, 242.

Tostegoptera lanceolata BLANCH. Cat. Col. Mus. Paris, 149; BURM. Lamell. II, 2d, 356.

Ancylonycha lanceolata LACORDAIRE, Gen. Coleopt., III, 285.

Throughout Kansas, extending into Eastern New Mexico and Upper Texas. The male is winged, while the female is apterous. The genus *Tostegoptera*, as very properly observed by Lacordaire (Gen. Col. III), does not differ from *Lachnosterna* by any essential characters.

DIPLLOTAXIS KIRBY.

D. obscura, oblonga nigro-pieca, subnitida, capite hemihexagono, dense punctato, antice late emarginato, margine anguste reflexo, sutura frontali profunda, vertice late bifoveato, thorace brevi dense punctato, medio paulo latiore, lateribus rotundatis, angulis anticis subaentis, posticis subrectis, margine laterali reflexo, ad angulos latiore impresso, clytris punctatis, tricostatis, tibiis anticis tridentatis. Long. .45.

One specimen from the Black Hills: Dr. Hammond. Resembles somewhat *D. brevicollis* Lec., but is more densely punctured, with the sides of the thorax more rounded, and more deeply impressed at the angles: the two foveæ of the vertex, if a constant character, will serve easily to distinguish it.

DIAZUS LEC.

Corpus oblongum alatum, breviter pubescens; caput mediocre, clypeo marginato, hemihexagono, sutura frontali parum distincta; labrum transversum late emarginatum; palpi maxillares articulo ultimo elongato-ovali, acuto haud impresso; mandibulae vix prominulae obtusae; mentum quadratum antice concavum; antennae breves, 9-articulatae, clava parva triphylla. Tibiae anticae unicalcaratae, 3-dentatae; posteriores paulo incrassatae, obsolete bicoronatae, calcaribus parvis; tarsi antici tibiis haud longiores, intermedii tibiis paulo longiores, postici tibiis breviores, crassiusculi; articulis omnium 4 primis aequalibus, 5to paulo longiore, unguiculis haud dentatis. Thorax linea apicali nulla, margine membranaceo distincto.

A genus belonging to the *Diplotaxes*, and having the characteristic form of abdomen of that tribe, viz., the 5th ventral segment united without suture to the propygidium, with the spiracle each side midway between the anterior and posterior margin. It differs, nevertheless, remarkably from the other genera by the nine-

jointed antennæ and simple unguis. The rough sculpture and short sparse pubescence give the insect very much the appearance of an *Ochodæus*.

D. rudis, ferrugineo-fuscus, oblongus, parum nitidus, pilis brevibus minus subtilibus, albidis minus dense vestitus, capite punctato, hemihexagono, clypeo antice latius emarginato, thorace convexo antrorsum angustato, lateribus valde rotundatis, dense punctato, elytris rugose punctatis, sutura costisque solitis tribus parum elevatis, pygidio confertim punctato. Long. .30.

Two specimens from the black hills: Dr. W. A. Hammond. The punctures are large and not deep. The body beneath is more shining, and less densely punctured.

OMORGUS Er.

O. scutellaris, apterus, niger, thorace cataphracto valde inæquali, lateribus rotundatis, elytris seriatim minus distincte punctatis, transversim cælati, interstitiis subquadratis, subæqualiter parum elevatis. Long. .6—65. Tab. I, fig. 4.

LEC. Proc. Acad. Nat. Sc. VII, 214.

Trox scutellaris SAY, Journ. Acad. Nat. Sc. III, 233.

From Platte river to Santa Fé. Other nearly allied species found in Texas are described by me in the place above cited; from all of them, however, it may be distinguished by the assemblage of characters here given.

CANTHON ILLIGER.

C. praticola, subovatus, ater opacus, supra subtilius licet evidenter granulatus, clypeo 6-dentato, dentibus duobus mediis obtusis, reliquis latis modice prominulis, thorace brevi convexo, lateribus valde rotundatis, elytris capite thoraceque vix longioribus, basi late emarginatis, lateribus et apice rotundatis, planiusculis versus scutellum paulo impressis, striis parum distinctis; thorace subtus versus angulum anticum tuberculo minuto marginali instructo, pro receptione pedum parum excavato, pygidio convexiusculo, parce subtiliter granulato. Long. .27—37.

Kansas and New Mexico. Resembles *C. nigricornis*, but the form is less abbreviated, the lateral teeth of the clypeus are more prominent, the elytra are less sparsely granulate, and the pygidium is more finely granulate, not impressed or carinate towards the base.

For the purpose of enabling the relations between the species of *Canthon* inhabiting the United States to be understood, as well as of providing names for several nondescript species, I add the following synoptic table:—

A. Prothorax pleuris linea transversa a coxis procedente obsoleta vel nulla.

a. Prothorax pleuris margine versus angulum anticum tuberculo minuto subinterrupto.

* Clypeo rotundato, bidentato; (species magnæ).

rotundatus supra granulatus, oculis latiusculis

vigilans.

rotundatus, oculis angustissimis;

supra evidenter granulatus

lævis.

obsolete granulatus

chalcites.

** Clypeo 6-dentato; (species minores).

§ Pygidio rude granulato, basi plus minus carinato;

clypei dentibus omnibus magnis;

oblongo-rotundatus, supra rude granulatus

ebenus.

rotundatus, supra subtilius granulatus

depressipennis.

clypei dentibus lateralibus minus prominulis;

breviter rotundatus, supra subtiliter, elytris parcius granulatus

nigricornis.

- §§ Pygidio vix grannlato, haud carinato ;
 ovato-rotundatus, supra granulatus *praticola.*
 rotundatus, supra haud granulatus *abrasus.*
- b. Prothorax pleuris margine versus angulum antieum tuberculo nullo (pro receptione pedum parum concavis).
- * Clypeo 6-dentato, dentibus externis parum prominulis ;
 rotundatus, ater, granulis depressis parvis obsitus *simplex.*
- ** Clypeo 4-dentato, dentibus internis acutis ;
 rotundatus, nitidus, nigro-cyaneus, punctulatus *cyanellus.*
- B. Prothorax pleuris linea transversa a coxis procedente notatis, antice pro receptione pedum subito declivibus ; (clypeus dentibus internis obtusis reliquis obsoletis).
 thorace punctato, elytris subopacis (cupreus, vel viridis) *viridis.*
 thorace fortius punctato, elytris nitidis punctulatis *perplexus.*

The species in the above synopsis, not before mentioned, are :—

C. depressipennis. A black, or greenish black species found in Georgia, resembling very closely *C. ebenus*, and only differing from that species by the body being less oblong, the thorax less suddenly rounded on the sides, and the granulations of the upper surface more minute. Long. .30—.35.

C. abrasus. A small rounded dull black species, from Georgia, with the eyes broader than usual, having no visible sculpture on the head and thorax, and very feebly granulate on the elytra, which are depressed in the scutellar region: the striæ are very indistinct, the pygidium not granulate. Long. .24.

C. cyanellus. A shining, very dark blue rounded species found in Texas, and given me by Mr. Ulke, having the clypeus oblique on the sides, and 4-dentate in front, the middle teeth being narrower than in the other species; the thorax is very minutely punctulate, much rounded and almost angulated on the sides, with a fovea remote from the margin opposite the widest part; the elytra are feebly punctulate, obsoletely striate, impressed at the scutellar region. Pygidium dull, obsoletely punctulate. Margin of the prothorax beneath without the small tubercle usually seen near the anterior angle. Long. .37.

Synonyms to be observed are :—

C. lævis: *Scarabæus lævis* Drury, Oliv.; *Scarabæus volvens* Fabr.; *Ateuchus volvens* Fabr.; *Scarabæus pilularius* † De Geer. *Coprobium obtusidens* Ziegler, is a slight variety of this species.

C. viridis: *Copris viridis* Beauv.; *Onthophagus viridicatus* Say. *Ateuchus obsoletus* Say, is a copper-colored variety of this species, which is more frequently found on leaves than in any other situation. Chevrolat (Ann. Ent. Soc. France, 2d ser. X, 632) refers Beauvois' species to *Onthophagus*, but I do not see anything in the figure or description to warrant such a reference.

Ateuchus probus Germ., Ins. Nov. 98, probably belongs to this genus, but I do not know any species that agrees with the description.

MELANOPHILA Esch.

M. miranda, nigra, ænescens, subtiliter dense punctulata, capite guttis quinque politis, circulo positis, thorace guttis quatuor anticis (intermediis maioribus), altera utrinque pone medium, lineaque dorsali postica nitidis levigatis; elytris margine basique tenuiter rubris, lineisque flavis decussatis ornatis. Long. .65. Tab. II, fig. 7.

LEC. Trans. Am. Phil. Soc. XI, 212.

Phænops mirandus LEC. Proc. Acad. Nat. Sc. VII, 83.

One specimen found at Fort Union, by Major Sibley. This is the most elegant species of the genus, the name of which seems by no means appropriate, yet being older than that of *Phænops* must be retained.

CHRYSOBOTHRIS ESCH.

C. quadrilineata, fusco-ænea, punctata opaca, latiuscula, capite pubescente, thorace lineis quatuor elevatis nitidis, medio valde canaliculato, elytris utrinque lineis elevatis nitidis, marginali suturalique integris, hac antice latiore, intermediis duabus confluentibus et pone medium interruptis. Long. .48—.60.

LEC. Trans. Am. Phil. Soc. XI, 233.

Santa Fé: Mr. Fendler. Broader and more robust than the other species found in the United States, and easily known by the sculpture of the thorax. The latter is twice as wide as its length, very deeply channelled in the middle, with a broad shining sparsely punctured entire vitta each side of the channel, and another each side which does not reach the apex; the depressed portions are opaque and very densely punctured. The elytra are wider than the thorax, serrate at the sides and apex, with an entire elevated shining line near the suture, dilated anteriorly so as to reach the suture, and an entire slender elevated line parallel with the outer margin; between these are two elevated lines which are confluent in two places, and interrupted behind the middle; the base is very deeply foveate as usual: the depressed portions are all densely punctured, and without lustre. Body beneath coarsely punctured, prosternum flat, covered with long white hair, with a slightly elevated smooth medial line.

C. cuprascens, obscure cupreo-ænea, longiuscula, minus depressa, fronte viridiæneo, dense punctato pubescente, thorace irregulariter punctato, vage impresso et subcanaliculato, elytris confertim punctatis, utrinque bimpressis, lineis nitidis elevatis, suturali marginalique integris, discoidali interna bis interrupta, externa ad medium postice abbreviata. Long. .31.

LEC. Trans. Am. Phil. Soc. XI, 234.

One specimen: Mr. Fendler. Resembles some of our smaller species, but is more convex and more coppery. The posterior impression of the elytra is deeper and larger than the anterior one, and although slightly lobed is not sinuous; the elevated lines are nearly smooth. The anterior tibiæ of the male are slightly bent, and armed with a tooth internally near the tip.

PSILOPTERA SOLIER.

P. Woodhousei, ænea, nitida, chalybeo-variegata, thorace confertim punctato, brevi, lateribus valde rotundatis, autrossum angustato, angulis posticis rectis, elytris apice integris, seriatim crenatis, maculis irregularibus opacis transversis profunde impressis. Long. .72—.97. Tab. II, fig. 6.

LEC. Trans. Am. Phil. Soc. XI, 195.

Dicrca? *Woodhousei* LEC. Proc. Acad. Nat. Sc. VI, 68.

Var. major, *Psiloptera valens* LEC. Proc. Acad. Nat. Sc., 1858, 66.

Creek Boundary Expedition: Dr. S. W. Woodhouse; Texas, Mr. Schott.

Body coppery bronze, varied with bluish reflections, moderately stout and convex; head strongly punctured, with three faint confluent elevated lines on the front;

labrum green; thorax cribrate punctate, more than twice as wide as broad as long, narrowed in front, very strongly rounded on the sides, narrowed a little towards the posterior angles, which are rectangular and sharp. Elytra with rows of large punctures, and with numerous deep impressed subconfluent spaces, which are opaque, densely punctured and finely pubescent; tip entire. Legs green bronzed, knees, tip of the tibiæ, and tarsi steel blue. Tip of the abdomen very slightly truncate in the male, rounded in the female.

CHAULIOGNATHUS HENTZ.

C. basalis, elongatus niger, thorace fulvo nitido, subrotundato, marginato, disco plus minus nigro, elytris rugose punctatis, flavo-fulvis macula magna communi triangulari basali, trienteque postico nigris, abdomine flavo, sæpe nigro annulato, articulo ultimo nigro. Long. .42.

Abundant near Fort Bridger and in the Black Hills: Dr. Hammond, Mr. Drexler. Varies much in the size of the black spot of the thorax, which is sometimes emarginate, and sometimes even divided into four dots. The antennæ of the male are as long as the body, of the female two-thirds as long. The thorax in *C. limbicollis* and *scutellaris* is opaque, while in the present it is very smooth and shining.

NIPTUS BOIELDIEU.

N. ventriculus, rufo-piceus, nitidus, pube sordida minus subtili dense vestitus, thorace globoso, grosse punctato, postice valde coarctato, tuberculis parum elevatis quatuor transversim positus, e pillis efformatis; elytris ovalibus, ventricosis, seriatim punctatis, setis crectis sat dense hispidis. Long. .10.

Santa Fé: Mr. Fendler. The elytra are twice as wide as the thorax, and about one-half longer than wide; the punctures are tolerably large and distant, and become smaller on the declivous posterior portion. The second joint of the antennæ is as long as the third.

TRYPOPITYS REDT.

T. punctatus, elongatus, fuscus sericeo-pubesceus, thorace ad basin utrinque transversim excavato, et in medio breviter earinato, elytris striis e punctis digestis vix impressis. Long. .25.

One specimen: Mr. Fendler. Precisely resembles in appearance *T. sericeus*, (*Xyletinus sericeus* Say, Journ. Acad. Nat. Sc. V, 171), but the thorax is less deeply excavated at base, and the striæ instead of being deep and cribrate are hardly impressed, and the interstices are flat.

PACTOSTOMA LEC.

P. anastomosis, piceus, pillis sordidis setiformibus parce vestitus, thorace punctato, subcanaliculato, latitudie brevior lateribus rotundatis marginatis, basi subtruncato, angulis posticis obtusis, elytris ovalibus postice subacutis et declivibus, parce punctatis, sutura, margine costisque tribus acute elevatis, quarum prima recta, secunda et marginali flexis cum prima confluentibus, tertia irregulari utrinque valde abbreviata. Long. .42—.5. Tab. II, fig. 11.

LEC. Journ. Acad. Nat. Sc. 2d Ser. IV, 19.

Microschatia anastomosis LEC. Proc. Acad. Nat. Sc. VI, 446.

Asida anastomosis SAY, Journ. Acad. Nat. Sc. 3.

Pelecyporus anastomosis LEC. Ann. Lye. Nat. Hist. V, 129.

Ologlyptus anastomosis LAC. Gen. Col. V, 159.

Kansas and New Mexico, near the Rocky mountains. The first dorsal costa is nearly straight, and extends from the base to the apex; the second commences near the base, and being curved, unites with the first about one-third from the tip; the third is irregular, running from the anterior third to the posterior fourth; the marginal one is entire, and unites with the first near the tip.

ASIDA LATR.

A. opaca, oblonga atra, opaca, brevissime parce flavo-pubescent, capite thoraceque scabro-punctatis, illo antice transversim excavato, hoc antroorsum angustato, latitudine brevior, lateribus rotundatis late subdepressis postice subsinuatis, angulis omnibus acutis, posticis paulo productis, elytris parce subtiliter minus profunde punctatis vage inaequalibus, nonnunquam obsolete sulcatis, lateribus marginatis, humeris obtusis. Long. .5—'68. Tab. I, fig. 9.

SAY, Journ. Acad. Nat. Sc. III, 254.

Euschides opaca LEC. Annals of the Lyceum of Nat. Hist. V, 127.

Kansas and New Mexico, near the mountains: specimens from the latter locality are smaller, and have the elytra much more uneven than those found near Platte river, but do not appear sufficiently distinct to be considered as another species.

EUSCHIDES LEC.

E. convexa, nigra subnitida, capite punctato antice transversim sulcato, thorace latitudine sesqui brevior, subtiliter parce punctato, lateribus rotundatis, margine punctato anguste reflexo, angulis anticis acutis, posticis obtusis haud rotundatis, elytris transversim convexis, pone basin thorace paulo latioribus, ad apicem valde declivibus, subtiliter rugosis lineis tribus obsolete utrinque notatis, humeris marginatis prominulis. Long. .85. Tab. I, fig. 10.

Arkansas river, near the mountains. Nearly allied to *Eu. obovata* Lee.; but with the thorax more convex and less broadly margined, and the elytra less obovate and more convex transversely. From *Eu. convexicollis* Lec. it differs by the larger and more strongly margined thorax.

PELECYPHORUS SOLIER.

P. sordidus, picco-niger, capite modice transversim impresso, punctato, thorace latitudine fere sesqui brevior ante medium angustato, lateribus rotundatis subserratis deplanatis et subreflexis, confertim grosse inaequaliter punctato, elytris oblongo-ovalibus, convexis postice subaeutis, sutura margine costisque utrinque duabus elevatis, lineis transversis sinuatis inter se irregulariter connexis; prosterno postice late sulcato. Long. .75—'85. Tab. I, fig. 11(♀).

LEC. Proc. Acad. Nat. Sc. VI, 446.

Arkansas river, near the mountains: also found in New Mexico, by Dr. Webb, of the Mexican Boundary Commission. In the female, the elytra are twice as wide as the thorax, and more rounded on the sides than in the male. The specimens are covered with a yellowish earthy substance which adheres closely.

EUSATTUS LEC.

E. reticulatus, rotundatus convexus, ater opaeus, thorace obsolete punctulato, lateribus marginatis paulo reflexis, angulis posticis valde productis, elytris foveis quadratis vagis impressis, obsolete sulcatis, tibiis anticis extrorsum serrulatis, versus apicem laevibus. Long. .46—'53. Tab. I, fig. 7 (♂, tibia antica).

LEC. Ann. Lyc. of Nat. Hist. V, 132.

Zophosis reticulata SAY, Journ. Acad. Nat. Sc. III, 250.

Found near the Rocky mountains from Platte river to Santa Fé, and westward as far as Tucson. Larger than *E. convexus* Lec., and less convex; the sides of the thorax are not suddenly depressed behind the middle as in that species.

EMBAPHION SAY.

E. contusum, atrum opacum, thorace latitudine fere duplo latiore, antice profunde emarginato, lateribus valde rotundatis, disco parce punctato, parum convexo, margine late explanato modice reflexo, angulis anticis subaentis, posticis latis obtusis valde rotundatis, ad basin medio reete truncato, elytris dorso planis, postice valde declivibus et acute angustatis, thorace vix latioribus, fortiter reflexo-marginatis, seriatim subtilius sat dense muricato punctatis, ad apicem singulatim breviter acuminatis (♀), vel in cauda brevi prolongatis (♂). Long. .55—.65. Tab. I, fig. 8.

Lec. Journ. Acad. Nat. Sc., Philad., 2d ser. IV, 40.

Fort Laramie and Santa Fé. Though differing very much in form from the *Helæus* like *E. muricatum* Say, the forms of the antennæ, oral organs and legs require it to be associated with that species. The figure given saves the necessity of a long description, while the diagnosis above will enable the species to be readily recognized.

Two specimens found by Dr. Webb, near the Mexican boundary, differ in having the sides of the thorax much more strongly reflexed, so that that part becomes much narrower than the elytra. This is, probably, merely a local variety, as similar differences occur among individuals of *Cyclurus clavatus*, and in many other insects which have the margins of the body widely reflexed.

BLAPSTINUS LATR.

B. pratensis, elongato-ovalis convexus, ater opacus, breviter subtiliter fusco-pubescent, capite thoraceque punctatissimis, hoc latitudine sesqui brevior, lateribus late rotundatis, ad apicem emarginato, angulis anticis acutis ad basin sinuato, angulis posticis rectis, elytris striis fortius punctatis, interstitiis paulo convexis, punctulatis. Long. .20—.23.

Platte River Valley, abundant. Differs from the Californian *B. pubescens* Lec., by the finer pubescence, by the more widely rounded sides of the thorax, and by the more prominent anterior angles.

B. vestitus, elongato-ovalis, convexus, æneo-niger, pube albida minus subtili dense vestitus, capite thoraceque confertim punctatis, hoc latitudine vix brevior, antrorsum paulo angustato, lateribus late rotundatis, ad apicem emarginato, angulis anticis acutis, ad basin sinuato, angulis posticis rectis, elytris striis punctatis, interstitiis planis punctulatis. Long. .20.

Two specimens from Platte River Valley. Very closely allied to the preceding, but the coarse pubescence, and the flattened intervals of the elytra seem to be sufficient characters for separating it.

CENTRONOPUS SOLLER.

C. opacus, elongatus, niger opacus, capite postice punctato, fovea verticali notato, antice concavo, epistomate late rotundato et fortiter marginato, thorace quadrato, antice vix angustiore, lateribus parum rotundatis, angulis posticis acutis, parum convexo, parce punctato, elytris striis e punctis parvis constitutis. Long. .57.

One male; Black Hills, Dr. Hammond. Differs from the Mexican *C. suppressus*, according to description, by being opaque, with finer elytral striae. The sexual

characters are nearly as in that species; the anterior femora have on the anterior face near the base an oval patch of yellow hair; the tibiæ are dilated internally at the middle into an obtuse angle, then broadly emarginate to the tip, at the inner side of which is a dense short brush of yellow hair; the first three joints of the anterior and middle tarsi are dilated, and furnished beneath with a dense brush of yellow hair; the middle tibiæ are thick internally at the apex and have also a brush of hair.

XYSTROPUS SOLIER.

X. pinguis, ater opacis, subtilissime punctulatus, brevissime pubescens, thoracæ brevi, lateribus valde rotundatis, basi sinuato, angulis posticis haud prolongatis, elytris striatis, interstitiis paulo convexis. Long. .37.

Santa Fé, Mr. Fendler: one specimen. Resembles *X. brevis* (*Cistela brevis* Say), but differs in the antennæ and feet being entirely black. The thorax is less deeply sinuate at the base, and the posterior angles are less acute, and not prolonged.

CYSTEODEMUS LEC.

C. vittatus, niger, capite thoracæque opacis, vage grosse punctatis, illo ad basin fere truncato, hoc subquadrato, subtiliter canaliculato, ad basin valde emarginato, elytris brevibus, inflatis obtusis, cribrato-punctatis, vitta angusta rubra abbreviata ornatis, abdomine amplissimo lævigato. Long. .5—·7. Tab. II, fig. 9.

LEC. Proc. Acad. Nat. Sc. VI, 330.

New Mexico, Dr. Wislizenus: found on the journey from Santa Fé to Chihuahua.

C. wislizeni, cyanescens, nitidus, capite thoracæque grosse punctatis, illo postice rotundato, hoc pentagono, canaliculato, angulis lateralibus acutis, elytris splendide cyanicis, sphericis, foveis profundis confertis cancellatis. Long. .45—·65. Tab. II, fig. 10.

LEC. Ann. Lye. Nat. Hist. of New York, V, 158; Proc. Acad. Nat. Sc. VI, 330.

Drs. Wislizenus and Webb, found in the same region with *C. vittatus*.

MORDELLA FABR.

M. insulata, longiuseula, nigra, capite thoracæque cinereo haud dense pubescentibus, hoc latitudine brevior, lateribus late rotundatis, elytris pube nigra indutis, sutura basique cinereo-marginatis, vitta obliqua a humero ad quadrantem extensa, maculaque transversa pone medium dense cinereo-pubescentibus; subtus maculis lateralibus cinereo-pubescentibus, antennis thoracæ longioribus basi, tibiis tarsisque anticis fusco-testaceis, stylo anali elongato. Long. .18—·21.

Two specimens, from Fort Riley; Mr. John Xantus. Approaches *M. oculata* Say, more nearly than any other species, but is readily distinguished by the markings as above described; the oblique vitta from the humerus does not reach the suture; the scutellum is densely clothed with cinereous hair.

MORDELLISTENA COSTA.

M. æmula, elongata, parallela, nigra, pilis pallide sericeis obsita, elytris evidenter punctulatis, pedibus posticis, tibiis carinulis numerosis brevibus obliquis, tarsorum articulo 1mo tibia vix brevior, 5-carinulato, 2d bicarinulato. Long. .18.

One specimen: Platte river. Elongate, not attenuated behind, black, above and beneath with gray silvery hair; head and thorax very finely punctulate, the latter

scarcely wider than long, somewhat narrowed in front, slightly rounded on the sides, bisinuate at the base, with the middle lobe broadly rounded. Elytra more than three times as long as the thorax, parallel on the sides, obtusely rounded behind, distinctly punctulate. Anal style slender. Posterior tibiae broad, with six or seven very short oblique ridges on the supero-external margin; first joint of posterior tarsi not shorter than the tibia, with five similar oblique ridges; second joint with two.

M. divisa, fusco-nigra, supra dense helvo-pubesceus, elongata, postice subattenuata, capite thoracque ante medium rufo-testaceis, elytris evidenter punctulatis; pedibus posticis, tibiis carinulis 2 vel 3 minutis obliquis, tarsorum articulo 1mo tibia haud brevior carinulis 4, 2do duabus instructo; antennarum basi pedibusque anterioribus rufo-testaceis, unguiculis simplicibus. Long. .11.

Platte river: resembles *M. marginalis* (*Mordella marg.* Say), but differs by the pubescence being coarser, more abundant, and of a yellowish brown color. The thorax is a little wider than long, not narrowed in front, broadly rounded on the sides, feebly bisinuate at base; the anterior half, with the head is reddish yellow, and the outline separating the yellow from the black is sinuate as in the species above mentioned.

In thus introducing the present genus into the fauna of the United States, a few remarks regarding our numerous species of *Mordellidae* will be appropriate. The genera which they represent are five in number, and may be thus separated:—

A. Scutellum quadratum, transversum postice submarginatum	<i>Tomoxia</i> COSTA.
B. Scutellum triangulare, apice rotundatum;	
a. Palpi maxillares articulo ultimo valde dilatato securiformi	<i>Glipa</i> LEC.
b. Palpi maxillares articulo ultimo triangulari;	
Antennae articulis externis valde transversis, velutinis	<i>Sphalera</i> LEC.
Antennae articulis externis triangularibus, haud transversis;	
Pedes postici simplices	<i>Mordella</i> LINN.
Tibiae tarsiue postice carinulis notati	<i>Mordellistena</i> COSTA.

To *Tomoxia* belong *M. bidentata* Say, and a nearly allied smaller species.

To *Glipa* belongs only *M. hilaris* Say. The body is slender; the outer joints of the antennae triangular; the maxillary palpi elongated, with the last joint in the form of an obtuse isosceles triangle, attached by its apex. The middle tibiae are longer than the tarsi; the posterior tibiae only moderately dilated, longer than the first joint of the tarsi, and destitute of oblique ridges; the anal style is short, truncate and submarginate.

Sphalera is the genus indicated, but not named by Lacordaire (Gen. Col. V, 609, note). The only native representative known to me is *M. melana* Germ.

To *Mordella* belong *Anaspis 4-punctata* Say; *M. 8-punctata* Fabr.; *oculata* Say; *insulata* Lec.; *seval* Say; *Anaspis triloba* Say; *M. marginata* Mels.; *lineata* Mels.; *scutellaris* Fabr. (*atrata* Mels.); *undulata* Mels., and four nondescript species.

To *Mordellistena* belong all the other species described by Say, Melsheimer, and myself; natural groups among them may be easily formed by the number and position of the tibial ridges, and the arrangement of the colors of the upper surface. *M. sericans* and one nondescript are remarkable by the posterior tibiae having a fine carina extending along the upper edge, and only a single very long oblique ridge



on the outer surface. These species do not indicate a generic separation, however, and represent the European *M. testacea*, the type of *Stenalia* Muls.

OPHRYASTES SCH.

O. vittatus, niger, albido-squamosus, rostro trisulcato, sulco medio profundissimo, lateralibus rectis profundis, mox ante oculos abbreviatis, thorace latitudine plus duplo breviorae inaequali, parce profunde punctato, lateribus sub-bituberosis, canaliculato, saepe fusco trivittato, elytris striis vix impressis, profunde punctatis, sutura cuprascente fusca, interstitiis 3io, 5to et 7mo fusco vittatis.

Long. .41—.48. Tab. I, fig. 13(♂). (*a.* rostrum).

SCHÖNHERR, Curc. 1,509; 5,819. LEC. Proc. Acad. Nat. Sc. VI, 443.

Liparus vittatus SAY, Journ. Acad. Nat. Sc. III, 316.

Platte and Arkansas rivers; found also at Eagle Pass, Texas. The elytra of the male are but little wider than the thorax; those of the female are more rounded, slightly flattened on the back, and about one half wider than the thorax. The dark vittæ of the thorax and elytra appear to be produced partly by abrasion, as specimens occur in which they are hardly to be seen; but the suture as far as the first stria is covered with brown scales having a coppery reflection.

CLEONUS SCH.

C. lutulentus, ater densissime sordide pubescens, vix variegatus, capite cum rostro grosse disperse punctato, hoc longitudinaliter vage impresso, medio vix carinato, thorace varioloso antrorsum angustato, lateribus antice rotundatis, disco paulo obscuriore, elytris inaequalibus, punctis magnis striatim positis. Long. .37—.43 (sine rostro).

Santa Fé, Mr. Fendler. Smaller and stouter than *C. trivittatus* Say, with the thorax more densely variolate, and hardly perceptibly excavated or carinated; the elytra have three or four vague impressions, not seen in that species.

C. pulvereus, alatus, niger sordidus, rostro carinato, cum capite vage grosse punctato, fronte foveata, thorace latitudine haud longiore, lateribus subparallels, antice breviter tubulato, angulis anticis rotundatis, grosse punctato ad basin medio late foveato, lateribus dense albo-pubescentibus, elytris convexis, oblongis postice obtuse rotundatis, striis fortiter punctatis haud impressis, totis dense albo-pubescentibus, lineis tribus curvatis obliquis sub-denudatis utrinque notatis; subtus albo-pubescentibus, obscure variegatus. Long. .5.

Arkansas river, one specimen. Of the same size and shape as *C. vittatus* Say, but very distinct by the thorax being not narrowed gradually in front, and not carinate, as well as by the different color.

C. angularis, niger, rostro carinato, cum capite fortiter vage punctato, fronte late foveata, thorace quadrato, lateribus subsinuatis, pone apicem valde constricto et ad apicem tubulato, angulis pone apicem rectis extantibus, varioloso-punctato, ad basin medio profunde foveato, elytris oblongis convexis, eribrato-striatis, interstitiis cinereo-pubescentibus spatiis denudatis variegatis, alternis cum sutura elevatis; subtus cinereo-pubescentibus, vix nebulosus. Long. .36—.45. Tab. I, fig. 12.

Two specimens: collected by Lieut. Beckwith. The first joint of the funiculus of the antennæ is slightly elongated, the others are short; the species, therefore, belongs to the division (called genus by Schönherr) Pachyeraerus. The projecting anterior angles of the thorax are very remarkable.

DEROBRACHUS SERVILLE.

D. geminatus, piceus, nitidus, thorace valde transverso, antice non angustato, parce punctulato, lateribus quadrispinoso, spina antica minore, elytris fere lavigatis, margine angustiore reflexo.

Long. 2·9. Tab. II, fig. 12 (femina).

Mas minor (long. 1·5—2·25), antennis longioribus crassioribus (fig. 12 a), pedibus anticis punctis elevatis exasperatis.

LEC. Proc. Acad. Nat. Sc. VI, 223.

New Mexico and Texas. The only female yet obtained was collected at Albuquerque, by Dr. T. C. Henry; several males were afterwards found by Messrs. Schott and Clark, of the Boundary Commission, at various places in Texas.

PRIONUS GEOFFROY.

P. curvatus, piceus, thorace subtiliter parce punctulato, spina anteriore tenui valde acuta, media hamata, angulis posticis subrectis prominulis, elytris subtiliter parce punctulatis et rugosis, spina suturali distincta; oculis magnis, antennis 12-articulatis. Long. 2·06.

One female from near Santa Fé. Resembles nearly *P. californicus* Motsch. (*crassicornis* Lec.), but the anterior spine of the thorax is more slender, and the middle one is curved backwards; the elytra are somewhat less punctured and rugose, but this is not a good character, as judging from a large series of *P. californicus*, it seems subject to variation.

P. fissicornis, nigro-piceus, nitidus, thorace subtiliter parce punctato, lateribus obtuse tridentato, elytris thorace latioribus parce sat grosse punctatis, obsolete costatis. Long. 1·0—1·6. Tab. I, fig. 14 (mas).

Mas antennis crassissimis valde imbricatis 27—30-articulatis, articulis subtus valde incisus (a).

Femina antennis tenuibus imbricatis, 25-articulatis, articulis subtus sinuatis (b).

HALDEMAN, Proc. Acad. Nat. Sc. III, 125; LEC. Journ. Acad. Nat. Sc. 2d ser. II, 108.

Platte river, New Mexico, and Texas, not rare.

P. emarginatus, piceus, nitidus, thorace punctatissimo, flavo-piloso, antrorsum angustato, lateribus ante medium unispinoso, angulis posticis rectis, elytris grosse parcius punctatis; antennis 14-articulatis. Long. ·75. Tab. II, fig. 13 (mas).

Mas antennis crassis valde imbricatis, articulis 3—13 ad apicem subtus valde emarginatis. Femina latet. SAY, Journ. Acad. Nat. Sc. III, 327; LEC. *ibid.* 2nd ser. II, 107.

Kansas and New Mexico, near the Rocky Mountains.

CRIOCEPHALUS MULS.

C. asperatus, piceus tenuiter pubescens, thorace subtransverso, lateribus obtuse angulatis, dorso plano ad latera granulis parvis exasperato, utrinque profunde lunatim excavato, medio subcanaliculato, ad basin fovea maiuscula impresso, elytris thorace parum latioribus, subtiliter rugosis, lineis utriusque duabus elevatis; oculis modice prominulis, genis subacutis; antennis corporis dimidio vix longioribus, versus basin crassiusculis, articulis ultimis quatuor subito brevioribus. Long. ·96.

Kansas and New Mexico. Differs from *C. agrestis* by the more flattened thorax, with more numerous elevated granules, and distinctly angulated sides. The antennæ are also heavier, and do not differ remarkably in length in the two sexes,

and their joints from the eighth to the eleventh are each one-half shorter than the seventh.

DRYOBIOUS LEC.

D. sexfasciatus, supra niger nitidus, capite thoraceque flavo-pubescentibus, illo inter antennis glabro transversim elevato, hoc fascia nigra ad medium ornato, latitudine paulo breviori, postice leviter coarctato, punctato, callis tribus dorsalibus notato, ante medium transversim impresso; elytris fasciis tribus apiceque late flavo-pubescentibus; subtus flavo-fasciatus, antennis pedibusque rufis, illis articulis internis subtus longe fimbriatis. Long. .65—1.0. Tab. I, fig. 15 (femina).

LEC. Journ. Acad. Nat. Sc. 2d ser. II, 23.

Callidium serfasciatum SAY, Journ. Acad. Nat. Sc. III, 415.

Alabama (Prof. Haldeman) and Ohio (Dr. Kirtland): Say found it on the Arkansas river. The antennæ of the female are but little longer than the body, those of the male are one-half longer. The femora are compressed and gradually slightly clavate.

ARHOPALUS SERV. (emend. LEC.)

A. charus, niger subtilissime punctulatus, thorace latitudine sesqui breviori, flavo-pubescente, fasciis tribus angustis nigris, elytris ad apicem oblique intus truncateis, a basi ad quadrantem flavis, gutta utrinque parva ad medium, fascia lata pone medium apiceque flavis, antennis pedibusque nigris. Long. .85. Tab. I, fig. 16 (var.).

LEC. Journ. Acad. Nat. Sc. 2d ser. I, 17; Proc. Acad. Nat. Sc. VI, 68.

Clytus charus SAY, Bost. Journ. Nat. Hist. I, 193.

Missouri, Dr. Engelmann; Creek Boundary, Dr. Woodhouse. The segments of the body beneath are edged with yellow; the humeri are marked with a black dot; the apical yellow spot sometimes includes a black spot, as in the specimen figured. I have seen only females of this species.

STENASPIS DUP.

S. solitaria, atra supra glabra, subtus parce cinereo-pubesceus, thorace latitudine fere duplo breviori, parce punctato, ante basin callo levi alteroque utrinque versus latera modice elevatis, lateribus ipsis bituberculatis, elytris subtilissime strigosis, parce subtiliter punctulatis, pedibus cyanescensibus. Long. 1.43. Tab. II, fig. 14 (femina).

LEC. Proc. Acad. Nat. Sc. VI, 441.

Cerambyx solitarius SAY, Journ. Acad. Nat. Sc. II, 410.

Callichroma solitarium HALD. Trans. Am. Phil. Soc. X, 32.

Smileceras solitarium LEC. Journ. Acad. Nat. Sc. 2d ser. II, 9.

New Mexico, Dr. Wislizenus; Kansas, Say; Texas, Messrs. Clark and Schott; Tucson, myself. The male has the antennæ longer than the body and slender.

TYLOSIS LEC.

T. maculatus, niger, thorace elytrisq. eoccineis, illo latitudine fere duplo breviori, punctato, callis quinque lævibus nigris, elytris confertim punctatis, macula humerali, alterisque utrinque tribus nigris. Long. .75. Tab. II, fig. 15.

LEC. Journ. Acad. Nat. Sc. 2d ser. II, 9.

New Mexico, Dr. Wislizenus (probably found west of Santa Fé). The humeral spot of the elytra in one specimen is obsolete. The antennæ of the male are much longer than the body.

ACMAEOPS LEC.

A. dorsalis, elongata, nigra opaca, supra parce breviter cinereo-pubesces, capite thoraceque confertissime punctatis, hoc latitudine subbreiore, convexo antice posticeque constricto, lateribus valde rotundatis, subcanaliculato, linea dorsali tenui levi, elytris thorace latioribus suparallelis, punctatis, vitta subsuturali lutea postice sensim attenuata, et pone medium abbreviata, apice rotundatis, subtus cinereo-pubesces, epimeris dense pubescentibus. Long. .45.

One specimen, from Republican Fork of Kansas river, kindly given me by Dr. T. B. Wilson. The antennæ are slender and reach two-thirds the length of the elytra; behind the middle of the elytra and extending nearly to the apex is a very obsolete yellowish stripe, which in other specimens may be found to be well developed, in which case there would be on each elytron a subsutural vitta abbreviated behind, and an external one abbreviated in front.

Resembles in appearance some of the varieties of *A. marginalis* Lec. (? *Leptura longicornis* Kirby), but the thorax is more robust and more rounded on the sides, the legs are entirely black, and the apex of the elytra more rounded.

LEPTURA LINN.

L. cribripennis, atra breviter parce pilosa, capite dense, thorace grosse punctato, hoc convexo, lateribus late rotundatis, utrinque constricto, elytris rubris subglabris, cribratim punctatis, ad apicem nigricantibus et intus oblique truncatis bidentatis, dente exteriori longiore; articulis antennarum 4, 5, 6, 9 et 10mo basi pallidis, 6to et 8vo pallidis macula apicali parva nigra notatis. Long. .7.

One specimen found by me on Platte river. Very nearly allied to *L. cinnamoptera* Kirby, but the elytra are much more coarsely punctured, and the lower half of the fourth joint of the antennæ is pale, while in the latter species there is merely a small pale spot at base.

MONILEMA SAY.

M. appressum, nigrum, thorace lateribus non spinosis, antice posticeque parce punctatis, elytris ante medium parce punctatis, dorso antice planis, lateribus subito valde declivibus. Long. .97. Tab. II, fig. 17.

LEC. Journ. Acad. Nat. Sc. 2d ser. II, 168.

New Mexico: one specimen: Dr. Wislizenus. Differs from the other species known to me by the flattened back and suddenly deflexed sides of the elytra.

STENOSTOLA MULS.

S. saturnina, plumbco-nigra, undique cinereo-pubesces, thorace latitudine haud longiore, punctato, lateribus subrotundato, pilis subtilibus erectis brevibus parce vestito, versus basin densius pubescente, elytris punctis remotis postice subtilioribus, ad apicem singulatim rotundatis. Long. .5.

One specimen from Kansas, kindly communicated to me by Mr. Burke. From its color it resembles at first sight *Superda moesta* and *concolor*, but the short convex front and dentate unguis will at once remind the student of the generic differences. Of the usual smooth thoracic spots seen in the species of this genus, only two very faint ones remain; they are situated about the middle, equidistant from each other and from the side.

AMPHIONYCHA LEC.

- A. ardens**, minus elongata cylindrica, atra, pube nigra erecta vestita, capite fronte vittisque duabus postice divergentibus late flammeo-pubescentibus, thorace supra flammeo-pubescente, vitta dorsali atra, latitudine sesqui brevior, lateribus parum sinuatis, elytris thorace paulo latioribus grosse punctatis, margine laterali ad humeros latiore, et ad dodrantem extenso ornatis. Long. .35.

Fort Riley: John Xantus, Esq. Broader than *A. flammata*, and readily distinguished by the thorax being one half broader than its length, with the lateral vittae much wider.

Also allied to these two is a third species found in Northern New York, and in New Hampshire; for specimens of it I am indebted to Mr. Henry Ulke, it may be thus characterized:—

- A. subarmata**, minus elongata cylindrica, atra, pube nigra erecta vestita, capite flammeo-pubescente, thorace latitudine vix brevior, lateribus sinuatis, medio tuberculo acuto promiulo armatis, vitta angusta sublaterali flammeo-pubescente, elytris grosse punctatis juxta suturam late sulcatis, sutura magis elevata. Long. .26—.28.

ÆDILIS SERV.

- Æ. spectabilis**, niger subtiliter dense cinereo fuscoque pubescens, thorace fusco maculis albis confluentibus utrinque ornato, elytris basi fasciisque tribus undulatis obliquis fuscis (intermedia latiore) notatis, pedibus fusco-annulatis, antennis longissimis cinereis, articulis 1—5 subtus dense pubescentibus, apice fuscis, 5to fasciulo apicali recurvato interno ornato. Long. .91. Tab. II, fig. 16.

LEC. Proc. Acad. Nat. Sc. of Philadelphia, VII, 82.

Fort Union: one specimen: Major Sibley. The figure renders unnecessary any farther description of this beautiful species.

LEMA FABR.

- L. trivirgata**, elongatula, parallela, testacea nitida, antennis (articulo primo excepto) thoracis maculis duabus (capitisque lateribus, abdominisque maculis sæpe), tibiæ apice tarsisque nigris; thorace latitudine longiore, medio valde coarctato, antice grossius parce punctato; elytris fortius punctato-striatis, interstitiis postice haud elevatis, vitta suturali alteraque submarginali angustis nigris. Long. .31.

Santa Fé: Mr. Fendler. Similar in appearance to *L. trilineata*, but known at once by the larger punctures of the thorax and elytra. The thorax is also more deeply constricted at the middle, and the outer vitta of the elytra is always narrower, usually extending only from the 9th to the 10th stria; in one specimen this vitta is interrupted behind the humerus. In another the second and third joints of the abdomen are marked each with a medial transverse black spot, and the occiput and two posterior spots of the head are black.

COSCINOPTERA LAC.

- C. franciscana**, oblonga, antice angustior, postice vix attenuata, nigra, subtus densius, supra sat dense albido-tomentosa, thorace latitudine sesqui brevior, lateribus obliquis parum rotundatis, punctato, linea angusta dorsali lævi, elytris fere dense punctatis, labro flavo-marginato. Long. .18—.20

Fort Riley: Mr. Xantus. Closely related to *C. dominicana*, but differs by the pubescence of the upper surface being much more dense, and by the elytra being more densely punctured. The sexual characters are as in that species.

CRYPTOCEPHALUS GEOFFR.

C. mucoreus, cylindricus, niger nitidus, subtus dense, supra subtiliter cinereo-pubescent, thorace convexo, subtiliter sat dense punctato, spatio parvo dorsali lævi, elytris basi margineque ultra medium late coccineis, maculaque coccinea rotundata apicali utrinque ornatis, punctis magnis striatim digestis, interstitiis parce subtiliter punctatis. Long. .25.

Fort Riley, one specimen: Mr. John Xantus. It has also been received from Texas by Mr. Ulke. A very distinct species; the elytral markings are a broad basal band curving along the margin to beyond the middle, and a small apical spot. Some of the varieties of *C. notatus* are similarly marked, and it is quite possible that the present species will be found also to vary in this respect.

PARIA LEC.

P. opacicollis, pallide flava, thorace latitudine fere duplo brevior, valde convexo, alutaceo subopaco, parce subtiliter punctulato, lateribus late rotundatis, elytris antice fortiter postice subtilius striatim punctatis, nitidis, margine postice, sutura, maculisque 4 sæpe nigris. Long. .13.

Fort Laramie: a very pretty species, readily distinguished by the faint sculpture of the thorax. The spots are placed, an oblong one at the middle of the base, and a larger quadrate one at the middle nearer the side than the suture.

P. punifa, saturate rufo-testacea, nitida, thorace convexo, quadrato, antice paulo angustato, lateribus parum rotundatis, parce punctato, elytris convexis, punctis paucis striatim positis ad basin majoribus, postice fere lævibus; antennis nigris, ad basin pedibusque flavis, pectore nigro. Long. .10.

One specimen sent me by the late A. Melly, Esq., as coming from Kansas. The striae consist only of a few distant large punctures towards the base, which become rapidly smaller and indistinct behind.

HETERASPIS.

H. nebulosa, cuprea, pube longa albida irregulariter vestita, capite inter oculos profunde angulatum impresso, fronte canaliculata, occipite bifoveato, thorace irregulariter punctato, latitudine haud longiore, ad basin truncato, lateribus rotundatis, elytris thorace latioribus convexis, striis grosse punctatis haud impressis postice obsolete, interstitiis alutaceis. Long. .15—.19.

Kansas, near Fort Laramie. The genus *Heteraspis* belongs in the vicinity of *Eumolpus*, and is distinguished from the other native genera by the following assemblage of characters:—

The mouth and eyes are not protected by the thorax, which beneath is simply truncate; the head is marked with a deep impression between the eyes; this impression is angulated each side, and continues around the inner posterior and inferior part of the eyes, forming a kind of orbit. The unguis is armed with a long acute tooth in *H. pubescens* Mels.; with a short acute tooth in *H. curtipennis* Mels., and in the next species, while in the one here described the tooth is almost obsolete.

H. smaragdula, viridi-anea, parce albo-pubesces, capite profunde angulatum impresso, fronte vix canaliculata, thorace latitudine haud longiore, lateribus rotundatis, ad basin late rotundato, punctato subrugoso versus apicem sublaevi, elytris longiusculis, convexis, thorace latioribus, striis haud impressis antice punctatis postice obsoletis, interstitiis alutaceis parce punctulatis. Long. ·20.

One specimen found at Fort Laramie.

MYOCHROUS.

M. squamosus, longiusculus, piceo-aneus, dense griseo-squamosus, breviter pubescens, capite thoraceque confertissime punctatis, hoc subrotundato, lateribus haud dentatis, elytris thorace non latioribus, punctis subquadratis striatum positis, postice obsoletis; tibiis anticis intus subarmatis, femoribus posticis haud dentatis; antennis versus basin rufis. Long. ·18.

Platte river, under dried buffalo excrement. Differs abundantly from *M. denticollis* by the absence of serration on the sides of the thorax, as well as by the posterior femora being not armed with a tooth; the scales of the upper surface of the body are broad.

The genus *Myochrous* consists of elongated pubescent *Enmolpi*, in which the thorax is slightly lobed behind the eyes; the latter are small and prominent; the antepectus is excavated for the reception of the head; the claws of the tarsi are not perceptibly toothed. The anterior tibiae are armed with a more or less prominent denticle on the inner surface near the middle; the posterior femora are sometimes simple, and sometimes toothed.

ÆDIONYCHIS LATR.

Æ. lugens, elongato-ovalis, atra opaca, thorace elytrisq. anguste marginatis, illo disperse punctato,, ad basin late rotundato, abdominis segmentis dorsalibus pallidis. Long. ·21.

Santa Fé, Mr. Fendler. Of the same shape, but smaller than *Æ. concinna* (*Ilaltica concinna* Fabr.), and easily known by its dull opaque black color, and by the base of the thorax not being sinuate near the posterior angles. The elytra appear impunctured; the pale color of the dorsal surface of the abdomen shows a little beneath at the margin and tip.

Æ. lobata, pallida, thorace angulis anticis acutis, lateribus late reflexis, elytris nigro-piceis subtiliter punctatis, basi versus scutellum latius, margine laterali trilobato, maculisque utrinque duabus disci pallidis, sutura anguste pallescente; subtus fusco-testacea, pedibus pallidis. Long. ·20.

One specimen from Kansas river. Allied to *Æ. quercata*, but larger, and with the elytra less broadly margined; the arrangement of the colors on the latter is also very different.

HALTICA ILLIGER.

H. punctigera, oblongo-ovalis, pallide flava, thorace latitudine duplo brevior, punctulato, lateribus rotundatis marginatis, dorso convexiusculo, punctis 4 nigris notato (externis sepe deficientibus), elytris dense subtiliter punctatis, sutura, vitta discoidea apicem haud attingente, alteraque juxta marginem nigris, tibiaram apicibus, tarsis antennisque fuscis, his articulis tribus primis subtus flavis. Long. ·30.

Kansas, near the Rocky mountains. Belongs with a numerous series of species allied to *H. alternata*; it is broader than that species, the thorax is more convex and more rounded on the sides; and the elytra are finely, but strongly and densely punctured.

II. pluriligata, pallide flava, elongata, thorace latitudine sesqui brevior, fere obsolete punctulato, guttis nigris quatuor, lineolaque dorsali versus basin ornato, lateribus rotundatis marginatis, elytris subtiliter parce punctulatis, sutura, vitta discoidea apicem haud attingente, alteraque juxta marginem nigris; tibiis ad apicem, tarsisque fuscis, antennis nigro-piceis, articulis tribus primis flavis supra infuscatis. Long. .32.

Kansas and Texas. Also allied to *II. alternata*, but narrower and with a less transverse thorax. Both this and the preceding, as well as the next species belong to Chevrolat's division (called genus by some authors) *Disonycha*.

II. cervicalis, ovalis, pallide flava, capite, antennis, pedibus, elytrisque nigris, his punctulatis subalutaceis; thorace subtilius marginato, levi. Long. .2.

Kansas and Georgia. Very similar to *II. collaris*, but rather stouter than that species, and distinguished by the uniform pale-yellow color of the under surface.

II. semicarbonata, ovalis convexa, fusco-atra, elytris opacis subtiliter punctatis, thorace parce punctulato, flavo, ad basin versus angulos posticos rectos sinuato, clypeo, abdominis segmento ultimo ventrali, coxis femoribusque flavis; antennarum articulis 1—3 subtus, tibiisque posticis ad basin flavis. Long. .24.

Santa Fé: one specimen: Mr. Fendler. Belongs to the division named *Disonycha* by Chevrolat, and resembles in appearance *II. collaris* Fabr., but is broader, having the form of *II. cervicalis*; it is readily distinguished from both by the dull elytra and different arrangement of the colors beneath.

II. ambiens, elongata, convexa, cyanea nitida, thorace virescente, punctulato, ad basin late rotundato, ante basin profunde transversim sulcato, sulco versus angulos profundiore, elytris subtiliter punctulatis, carina submarginali distincta, postice curvata et versus suturam oblitterata. Long. .22.

Santa Fé: Mr. Fendler. Differs from *II. subplicata* in the smaller size, and by the more elevated submarginal costa of the elytra being curved behind. It nearly resembles a nondescript species from New Hampshire (*II. alni* Harris), but the thorax is more convex, and the elytra more deeply punctured, while the curved portion of the costa is less prolonged towards the suture; in both the elytra are broadly sulcate about the middle, just within the costa.

II. subplicata, elongata, obscure cyanea, parum nitida, thorace latitudine paulo brevior, subtiliter punctulato, lateribus rectis angulis omnibus distinctis, postice fortiter transversim sulcato, elytris parce punctulatis, plica submarginali minus elevata versus apicem sensim oblitterata notatis. Long. .25.

One specimen: found in Platte river valley. Belongs to a numerous group of the division *Graptozona* Cherv., having an elevated fold parallel with the outer margin of the elytra; this fold in the other species is, however, much more elevated and curved posteriorly towards the suture.

II. punctipennis, elongata, læte viridi-ænea, thorace aurtorsum angustato, latitudine sesqui brevior, lateribus late rotundatis, vix subtiliter punctulato, postice transversim tenuiter obsolete sulcato, elytris thorace paulo latioribus, distincte licet subtiliter punctatis, antennis nigris. Long. .2.

Kansas and Santa Fé. Specimens from New Mexico were collected by the late Richard C. Kern, Esq., and were given to me by Prof. S. S. Haldeman. This is a *Graptozona* of slender form, and is readily distinguished by the distinctly punctured elytra, and by the slight posterior impression of the thorax.

II. obliterata, elongata, chalybeo-atra subopaca, thorace subtiliter punctulato, ad basin late rotundato, ante basin stria transversa subtili impresso, elytris subtiliter fere obsolete punctulatis. Long. .27.

One specimen, Mr. Fendler. Very distinct from other species of *Graptodera* known to me, by the fine transverse line of the thorax being obliterated towards the sides.

II. torquata, elongata, thorace punctulato ad basin late rotundato, ante basin profunde transversim sulcato, plus minusve cupreo, elytris chalybeis purpureo micantibus, confertim punctulatis, margine laterali cupreo-æneis; subtus obscure ænea, pedibus purpureo-chalybeis. Long. .2.

Santa Fé, Messrs. Fendler and Kern. An elongate *Graptodera* distinguished by the color and the fine dense punctuation of the elytra, from all the species that resemble it.

II. biteniata, elongata nigro-picea, supra permitida, capite postice pallidiore, thorace fusco-testaceo, maculis tribus obscuris, parce punctulato, ante basin transversim leviter impresso, et in medio foveato, elytris punctulatis nigris, vitta lata dorsali margineque (mox ante apicem sæpe connexis) pallidis; antennis pedibusque fusco-testaceis, femoribus posticis obscuris. Long. .18—.21.

Santa Fé, Mr. Fendler. Larger than *II. elongata* Fabr. (*teniata* Say), and with it, belongs to the division *Systema* Chevr. The marginal and dorsal vittæ sometimes unite near the tip.

LONGITARSUS LATR.

I. nigripalpis, elongata, alata, flavo-testacea, oculis palpisque nigris, fronte carinata, vertice tuberculis duobus parum elevatis, linea impressa definitis, thorace latitudine breviori, lateribus rotundatis, valde convexo, alutaceo, ante basin transversim sulcato, elytris thorace latioribus oblongis, subtiliter punctulatis pone basin late impressis; tibiis posticis elongatis haud sulcatis, calcari parvo terminatis. Long. .12.

One specimen, Platte river. The first joint of the hind tarsi is somewhat less than half the length of the tibia; the antennæ are lost. Differs from the other species by the non-sulcate hind tibiæ.

I. subrufus, rufo-testaceus, nitidus, fronte carinata, thorace latitudine haud breviori, angulis rotundatis, convexo, haud dense punctato, elytris thorace latioribus, ovalibus convexis, subtiliter punctatis, introrsum vix conspicue striatis; tibiis posticis sulcatis, calcare brevi terminatis, antennis corpore vix brevioribus, extrorsum infuscatis. Long. .12.

Fort Laramie. The upper edge of the hind tibiæ is finely serrate; the tarsus is as long as the tibia.

I. rubidus, apterus, ovatus, piceo-ferrugineus, nitidus, capite infuscato, fronte carinata, vertice transversim impresso, thorace transverso, basi et angulis rotundatis, convexo, alutaceo, subtiliter parce punctulato, elytris ovalibus convexis, sat dense minus subtiliter punctatis, antennis testaceis, extrorsum vix infuscatis, corpore paulo brevioribus. Long. .11.

Fort Laramie. The spur of the hind tibia is short; the tarsus is as long as the tibia; the elytra are obtusely rounded at tip, and the pygidium is slightly prominent.

GLYPTINA LEC.

G. spuria, testacea, nitida, fronte elevata, vertice bituberculato utrinque oblique impresso, thorace transverso, basi rotundato, lateribus parallelis angulis anticis oblique truncatis, punctato, elytris oblongo-ovalibus thorace latioribus, fortiter punctato-striatis; antennis extrorsum fuscis. Long. .08.

Fort Laramie. I have given the name *Glyptina* to a group of *Haltica* intermediate between genuine *Haltica* and *Longitarsus*. The head is marked with a deep oblique line each side between the eyes, running forward to the frontal elevation: the antennæ are half as long as the body, slightly thickened externally; the 2d and 3d joints equal in length: the thorax is quadrate, transverse, with the anterior angles obliquely truncate, the disc moderately convex, without impressed lines. The elytra oblong, wider than the thorax, with rows of punctures; the hind tibiæ deeply sulcate above, with the edge of the groove very finely serrate, the terminal spur small, the tarsi attached at the end of the tibiæ, 1st joint equal to the others combined, and nearly half as long as the tibiæ. Besides the two species here described, the Californian *Haltica ccrina* Lec., Pacific R. R. Expl. and Surveys, vol. xi, insects, 68, also belongs to this genus.

G. hissorques, testacea, nitida, fronte parum elevata, vertice utrinque oblique sulcato, thorace transverso, basi rotundato, lateribus parallelis, angulis anticis oblique truncatis, nitido, levi, vix obsolete parce punctulato, elytris oblongo-ovalibus, thorace latioribus, fortiter punctato-striatis; antennis extrorsum vix infuscatis. Long. .08.

One specimen with the preceding, which it entirely resembles, except in the sculpture of the head and thorax. It is also found in Pennsylvania.

CHLETOCNEMA STEPHENS.

C. subviridis, supra viridi-ænea, obscura, subnitida, thorace latitudine fere duplo brevior, lateribus obliquis late rotundatis, fortius sat dense punctato, elytris fortiter striato-punctatis, interstitiis subtiliter uniseriatim punctulatis, pedibus antennisque nigris, his basi testaceis, tibiis basi vix piecescentibus. Long. .12.

Fort Laramie. Broader than *C. denticulata*, and differing therefrom in many particulars. I have retained the older name for this genus, as I know no reason for the substitution of Redtenbacher's *Plectroscelis*, which is adopted by many naturalists of Germany and France.

LUPERUS GEOFFROY.

L. rufipes, elongatus, chalybeo-niger, thorace parce punctulato, latitudine hand brevior, lateribus late rotundatis subsinuatis, angulis posticis prominulis, elytris fere obsolete punctulatis, chalybeis, antennis pedibusque rufo-testaceis, illis extrorsum fuscis. Long. .22.

Santa Fé, Messrs. Kern and Fendler. Of the form and size of *L. meraca* (*Galleruca meraca* Say), but known by the uniform color of the feet, the less distinctly punctulate elytra, and the less rounded sides of the thorax.

MICRORHOPALA CHEVR.

M. lætula, elongata, postice modice dilatata, et obtuse rotundata, nigra, capite pone antennis, thoraceque toto late rufo-flavis, hoc parce cribrato, transverso, lateribus obliquis rectis, elytris punctis magnis striatis, seriebus per paria approximatis, internis postice subtilioribus, vittaque utrinque basali ante medium abbreviata flava. Long. .25.

Fort Riley, Mr. Xantus. Closely related to *M. vittata*, but differs by the sides of the thorax being entirely straight, as well as by the colors; the thorax is very narrowly margined with black, but is yellow both above and beneath, as is also the

head, excepting the antennæ, front and mouth, which are black. The outer intervals of the elytra are not shining, and are alternately a little more convex, as in *M. vittata*, and the edge is not serrate.

CASSIDA LINN.

C. ellipsis, elliptica convexa, supra testacea pallida, nitida (quando viva, læte aurea), thoracæ antice rotundato, margine late explanato hyalino, angulis lateralibus paulo rotundatis, elytris thoracæ haud latioribus, fortius striato-punctatis, margine subhyalino, antice declivi postice explanato angustius reflexo, capite corporeque nigris, abdomine anguste testaceo marginato, pedibus antennisque flavis, his articulis 3—6 tenuibus æqualibus, 7—11 latioribus, ultimis quatuor nigris. Long. .26.

One specimen, found near Long's Peak. Allied by the declivous and scarcely explanate but narrowly reflexed elytral margin, to *C. 6-punctata* and *C. 9-maculata*, but differing by the more narrow, regularly elliptical form of body. The legs in *C. 9-maculata* are black; in *C. 6-punctata* they are reddish yellow, but the punctures of the striæ of the elytra in that species are less approximate, each elytron is marked with three black spots, the humeral angles are more prolonged, and the side angles of the thorax are more rounded. Still more essential differences exist in the antennæ: in the present species the 2d joint is a little longer than wide; the 3d is slender, one-half longer than the 2d; the 4th, 5th, and 6th are equal in length and thickness to the 3d; the 7th is more than twice as wide, triangular, pale yellow, like the preceding joints; the remaining four joints are as wide as the 7th, and black. The middle of the anterior margin of the thorax is slightly emarginate, but I suspect this character to be accidental.

C. atripes, ovata, convexa, supra testacea (quando viva læte aurea), thoracæ antice rotundato, margine hyalino late explanato, angulis externis haud rotundatis, elytris thoracæ vix latioribus, humeris paulo productis rotundatis, striis e punctis parvis constitutis, margine hyalino antice declivi postice explanato, vix reflexo, utrinque guttis tribus nigris notatis; subtus nigra, abdomine testaceo-marginato, antennis basi flavis, articulis 3—5 æqualibus, 6to paulo latiore, ultimis 5 nigris. Long. .23.

One specimen, found near Long's Peak. Differs from *C. nigripes* chiefly by the thorax being nearly as wide as the elytra, with the lateral angles scarcely rounded: the margin of the elytra anteriorly is more obliquely declivous than in that species.

With regard to the position of *C. unipunctata* Say, I am somewhat doubtful, although to avoid the multiplication of genera it should probably be allowed to remain in *Cassida*. The 2d joint of the antennæ is scarcely thicker than the 3d, which is about twice as long; the 4th is two-thirds as long as the 3d, and somewhat wider; the 5th is as wide as its length; and the remaining ones are a little wider than their length: the last is oval and pointed. The unguis, as in our other species, are simple.

BRACHIACANTHA MULS.

B. tau, rotundato-ovalis, convexa, supra rufo-flava, nitida dense subtiliter punctulata, thoracæ basi subinfusato, elytris fascia basali ad humerum abbreviata, sutura anguste, guttisque utrinque duabus paulo pone medium nigris, margine externo pone maculas, apicalique nigro, subtus nigra, antennis, palpis, pleuris pedibusque flavis. Long. .19.

One specimen, Fort Riley: John Xantus, Esq. The head and sides of the thorax are paler than the rest of the upper surface.

EROTYLUS FABR.

E. Boisduvalii, oblongus, ater, subnitidus, elytris sat convexis, albido-testaceis, punctis maioribus nigris minus crebre impressis, singulo macula parva laterali nigra ornato. Long. 6—62.
Tab. II, fig. 8.

CHEVROLAT, Col. Mex. 2d cent.; Lac. Mon. Erotyl. 466.

A large number of this species was contained in the collection made by Mr. Fendler, near Santa Fé.



INDEX OF GENERA.

- | | | |
|--|---|--|
| <p> <i>Acmæops</i>, 21.
 <i>Ædilis</i>, 22.
 <i>Agabus</i>, 4.
 <i>Amblychila</i>, 1.
 <i>Amphionycha</i>, 22.
 <i>Anisodactylus</i>, 2.
 <i>Anisomera</i>, 5.
 <i>Arhopalus</i>, 20.
 <i>Asida</i>, 14.

 <i>Blapstinus</i>, 15.
 <i>Brachiacantha</i>, 28.

 <i>Calosoma</i>, 3.
 <i>Canthon</i>, 10.
 <i>Carpophilus</i>, 6.
 <i>Cassida</i>, 28.
 <i>Centronopus</i>, 15.
 <i>Chaetocnema</i>, 27.
 <i>Chauliognathus</i>, 13.
 <i>Chrysobothris</i>, 12.
 <i>Cleonus</i>, 18.
 <i>Coscinoptera</i>, 22.
 <i>Criocephalus</i>, 19.
 <i>Cryptocephalus</i>, 23.
 <i>Cymindis</i>, 2.
 <i>Cysteodemus</i>, 16.

 <i>Derobrachus</i>, 19.
 <i>Diazus</i>, 9.
 <i>Dicelus</i>, 3. </p> | <p> <i>Diplotaxis</i>, 9.
 <i>Dryobius</i>, 20.

 <i>Embaphion</i>, 15.
 <i>Erotylus</i>, 29.
 <i>Eusattus</i>, 14.
 <i>Euschides</i>, 14.

 <i>Glipa</i>, 17.
 <i>Glyptina</i>, 26.

 <i>Haltica</i>, 24.
 <i>Harpalus</i>, 2.
 <i>Heteraspis</i>, 23.
 <i>Hister</i>, 7.

 <i>Ilybius</i>, 4.

 <i>Lachnosterna</i>, 9.
 <i>Lema</i>, 22.
 <i>Leptura</i>, 21.
 <i>Longitarsus</i>, 26.
 <i>Luperus</i>, 27.

 <i>Melanophila</i>, 11.
 <i>Meligethes</i>, 6.
 <i>Micrixys</i>, 1.
 <i>Microrhopala</i>, 27.
 <i>Microschatia</i>, 13.
 <i>Monilema</i>, 21.
 <i>Mordella</i>, 16. </p> | <p> <i>Mordellistena</i>, 16.
 <i>Myochrous</i>, 24.

 <i>Niptus</i>, 13.
 <i>Nomaretus</i>, 3.

 <i>Edionychis</i>, 24.
 <i>Ologlyptus</i>, 13.
 <i>Omorgus</i>, 10.
 <i>Ophryastes</i>, 18.

 <i>Pactostoma</i>, 13.
 <i>Paria</i>, 23.
 <i>Pelecyphorus</i>, 14.
 <i>Phileurus</i>, 8.
 <i>Polyphylla</i>, 8.
 <i>Prionus</i>, 19.
 <i>Psiloptera</i>, 12.

 <i>Saprinus</i>, 7.
 <i>Silpha</i>, 6.
 <i>Sphalera</i>, 17.
 <i>Stenaspis</i>, 20.
 <i>Stenostola</i>, 21.

 <i>Thyce</i>, 8.
 <i>Tomoxia</i>, 17.
 <i>Trypopytus</i>, 13.
 <i>Tylosis</i>, 20.
 <i>Xystropus</i>, 16. </p> |
|--|---|--|

LIST OF SPECIES.

I. OF KANSAS AND NEBRASKA.

CICINDELIDÆ.

Amblychila Say.

- cylindricornis *Say*.
A. Piccolomini Reiche.

Megacephala Latr.

- virginica *Dej*.
Cicindela virginica Linn.

Cicindela Linn.

- pulehra *Say*.
 seutellaris *Say*.
 sexguttata *Fabr*.
 purpurea *Oliv*. var. *Auduboni* Lec.
 obsoleta *Say*.
 prasina *Lec*. Trans. Am. Phil. Soc. XI, 31.
 nigrocerulea *Lec*.
 cinetipennis *Lec*.
 pusilla *Say*.
 cyanella *Lec*. Trans. Am. Phil. Soc. XI, 46.
 terriola *Say*.
 cuprasens *Lec*.
 sperata *Lec*. Trans. Am. Phil. Soc. XI, 50.
 lepida *Dej*.
 hirticollis *Say*.
C. albohirta Dej.
 vulgaris *Say*.
C. obliquata Dej.
 fulgida *Say*.
 venusta *Lec*.
 formosa *Say*.
 punctulata *Fabr*.
 cumatilis *Lec*.
C. Guerci Chevr.
 circumpieta *Ferté*.
C. Johnsoni Fitch, N. Y. Agr. Soc. 1856, 487.
 celeripes *Lec*.
 cursitans *Lec*. Trans. Am. Phil. Soc. XI, 60.
 limbata *Say* (fide *Say*).
 decemguttata *Say* (fide *Say*).

CARABIDÆ.

Brachinus Weber.

- cyanipennis *Say*.
 and four other species.

Helluomorpha Lap.

- laticornis *Lap*.
Helluo laticornis Dej.
 praeusta *Lap*.
Helluo praeustus Dej.

Galerita Fabr.

- dubia *Lec*.
 'Carabus bicolor Drury,' fide Klug.
 atripes *Lec*. Proc. Acad. Nat. Sc. 1858, 59.

Lachnophorus Dej.

- elegantulus *Mann*.
Tachypus mediosignatus Ménét.

Casnonia Latr.

- pennsylvanica *Dej*.
Attelabus pennsylvanicus Linn.

Leptotrachelus Latr.

- dorsalis *Latr*.
Odacantha dorsalis Fabr.
Sphaeracra dorsalis Say.

Lebia Latr.

- viridipennis *Dej*.
L. borea Hentz.
 smaragdula *Dej*.
 viridis *Say; Dej*.
 pumila *Dej*.
L. floricola Harris.
 solea *Hentz*.
L. scapularis Dej.
 furecata *Lec*.
 axillaris *Dej*.

Blechnus Motsch.

- linearis *Schaum*. Ins. Deutschl. I, 275.
Dromius angustus||*Lec.*
Bomius linearis *Lec.*

Axinopalpus *Lec.*

- biplagiatus *Lec.*
Dromius biplagiatus *Dej.*

Glycia *Chaud.*

- viridicollis *Lec.*
Cymindis viridicollis *Lec.*
 purpurca *Lec.*
Cymindis purpurea *Say.*
 var. *Cymindis amocna* *Lec.*

Calleida *Dej.*

- punctata *Lec.*
 smaragdina *Dej.*

Cymindis *Latr.*

- laticollis *Say.*
 eribricollis *Dej.*
 cribrata *Lec.*
 pilosa *Say*
C. pubescens *Dej.*

Calathus *Bon.*

- gregarius *Dej.*
Feronia gregaria *Say.*

Rhadine *Lec.*

- larvalis *Lec.*

Platynus *Bon. emend. Brullé.*

- extensicollis *Lec. Proc. Acad. Nat. Sc. VII, 46.*
Feronia extensicollis *Say.*
Feronia (Anchomenus) proximus *Harris.*
Anchomenus extensicollis *Dej.*
Anchomenus obscuratus *Chaud.*
Anchomenus Lecontei *Lec.*
Anchomenus elongatulus†*Lec.*
Anchomenus viridis *Lec.*
 punctiformis *Lec. ibid. VII, 50.*
Feronia punctiformis *Say.*
Agonum rufipes *Dej.*
Agonum foveicollis *Chaud.*
 errans *Lec. ibid. VII, 50.*
Feronia errans *Say.*
 subcordatus *Lec. ibid. VII, 51.*
Agonum erythropum||*Kirby.*
 basalis *Lec. Proc. Acad. Nat. Sc. VII, 52.*
Agonum basale *Lec.*

- nutans *Lec. ibid. VII, 52.*

- Feronia nutans* *Say.*
Agonum femoratum *Dej.*
 picipennis *Lec. ibid. VII, 53.*
Agonum picipenne *Kirby.*
 chalceus *Lec. Agassiz' Lake Sup. 205.*
Agonum chalceum *Lec.*
 maculifrons *Say (Feronia), (fide Say).*
 scutellaris *Say (Feronia), (fide Say).*
 ? *Agonum melanarium* *Dej.*
 ? *Platynus melanarius* *Lec.*

Evarthrus *Lec.*

- sigillatus *Lec. Journ. Acad. Nat. Sc. 2d ser. II, 228.*
Feronia sigillata *Say.*
Feronia vidua *Dej.*
 seximpressus *Lec. ibid.*
Feronia seximpressa *Lec.*
 corax *Lec. ibid. II, 229.*
Feronia corax *Lec.*
 abdominalis *Lec. ibid. II, 232.*
Feronia abdominalis *Lec.*
 lixa *Lec. ibid.*
Feronia lixa *Lec.*
 incisus *Lec. ibid.*
Feronia incisa *Lec.*
 ovipennis *Lec. ibid.*
Feronia ovipennis *Lec.*
 latebrosus *Lec. ibid. II, 233.*
 constrictus *Lec. ibid.*
Feronia constricta *Say*
 substriatus *Lec. ibid.*
Feronia substriata *Say.*
 colossus *Lec. ibid.*
Feronia colossus *Say.*
 heros *Hald. Proc. Acad. Nat. Sc. VI, 361.*
Feronia heros *Say.*

Pterostichus *Bon.*

- caudicalis *Lec.*
Feronia caudicalis! *Say.*
Feronia luctuosa *Dej.*
Omascus nigrita†*Kirby.*
Pterostichus luctuosus *Lec. loc. cit. II, 243.*
 mutus *Lec.*
Feronia muta *Say.*
Feronia morosa *Dej.*
Omascus picicornis *Kirby.*

Pocilus *Bon.*

- scitulus *Lec.*
 cyaneus *Lec.*
 chalcites ———.
Feronia chalcites *Say.*
Pocilus Sayi *Brullé.*

lucublandus Kirby.

Feronia lucublanda Say.
bicolor Lec.

Amara Bon.

laticollis Lec.

? *Curtonotus convexiusculus* † Kirby.

carinata Lec.

furtiva Say.

libera Lec. Proc. Acad. Nat. Sc. VII, 349.

Isopleurus hyperboreus † Lec.

? *Curtonotus latior* Kirby.

angustata Say.

A. indistincta Hald.

impuncticollis Say.

A. difficilis Lec.

subpunctata Lec. Proc. Acad. Nat. Sc. VII, 352.

confusa Lec.

polita Lec.

obesa Say.

A. patricia † Dej.

Feronia obesa Say.

diffinis Lec. Proc. Acad. Nat. Sc. VII, 354.

Percosia diffinis Lec.

terrestris Lec. Proc. Acad. Nat. Sc. VII, 354.

Isopleurus terrestris Lec.

musculus Say.

Aerodon musculus et contempta Lec.

Nothopus Lec.

zabroides Lec.

Euryderus || *zabroides* Lec.

Geopinus Lec.

incrassatus Lec.

Daptus incrassatus Dej.

Cratognathus Dej.

setosus Lec. Trans. Am. Phil. Soc. X, 381.

Piosoma setosum Lec.

Cratacanthus Dej.

dubius Lec.

Harpalus dubius Beauv.

Cratacanthus pennsylvanicus Dej.

Agonoderus Dej.

lineola Dej.

Carabus lincola Fabr.

dorsalis Lec.

pallipes Dej.

Carabus pallipes Fabr.

Discoderus Lec.

parallelus Lec. Trans. Am. Phil. Soc. X, 382.

Selenophorus parallelus Hald.

tenebrosus Lec. *ibid.* 382.

Selenophorus tenebrosus Lec.

Spongopus Lec.

verticalis Lec.

Anisodactylus Dej.

rusticus Dej.

Harpalus rusticus Say.

Anisodactylus pinguis Lec.

Anisodactylus gravidus Lec.

Anisodactylus crassus Lec.

agricola Dej.

Harpalus agricolus Say.

Harpalus paradoxus Hald.

Anisodactylus striatus Lec.

baltimorensis Dej.

Harpalus baltimorensis Say.

cænus Dej.

Harpalus cænus Say.

Eurytrichus Lec.

terminatus Lec.

Feronia terminata Say.

Harpalus terminatus Dej.

Harpalus Latr.

impotens Lec. Journ. Acad. Nat. Sc. 2d ser. IV, 14.

pedicularius Lec.

Selenophorus pedicularius Dej.

troglydites Lec.

Selenophorus troglydites Dej.

æreus Lec.

Selenophorus æreus Lec.

Selenophorus planipennis Lec.

ellipticus Lec.

Selenophorus ellipticus Dej.

caliginosus Say.

Carabus caliginosus Fabr.

amputatus Say.

H. Stephensii Kirby.

rotundicollis Kirby.

pennsylvanicus Lec.

Carabus pennsylvanicus De Geer.

Carabus bicolor Fabr.

Harpalus faunus † Dej.

compar Lec.

H. pennsylvanicus Say.

H. bicolor † Dej.

stupidus Lec., ante, 3.

oblitus Lec., ante, 2.

nitidulus Chaud.

herbivagus Say.

ellipsis Lec.

ventralis Lec.

funestus Lec.

Stenolophus Dej.

- ochropezus Dej.
Feronia ochropeza Say.
 humilis Lec.
Acupalpus humilis Dej.
 dissimilis Dej.

Bradycellus Er.

- obesulus Lec.
 badiipennis Lec.
Geobænus rufierus † Lec.
Stenolophus badiipennis Hald.
 congener Lec.
Geobænus congener Lec.
 rupestris Lec.
Trechus rupestris Say.
Acupalpus elongatulus Dej.
Trechus flavipes Kirby.

Badister Clairv.

- notatus Hald.
B. terminalis Lec.
 micans Lec.

Diploohila Brullé.

- laticollis Lec.
 gens maior Lec.
 obtusa Lec.

Dicælus Bon.

- lævigatus Lec.
 splendidus Say.
 sculptilis Say.
 simplex Dej.
 var. *D. obscurus* Lec.
 elongatus Dej.

Oodes Bon.

- amaroides Dej.

Atranus Lec.

- pubescens Lec.
Anchomenus pubescens Dej.
Anchomenus obconicus Hald.

Chlænienus Bon.

- purpuricollis Randall.
 tomentosus Dej.
Epomis tomentosus Say.
Amara luctuosa Germ.
 pennsylvanicus Say.
C. vicinus Dej.
C. pubescens Harris.
 vafer Lec. Proc. Acad. Nat. Sc. VI, 66.

æstivus Say.

- C. cobaltinus* Dej.
 erythropus Germ.
C. rufilabris Dej.
 laticollis Say.

C. diffinis Chaud. Bull. Mosc. 1856, II, 279. *

lithophilus Say.

- C. viridanus* Dej.
 sericeus Say.
Carabus sericeus Forster.
Chlænienus perrivirdis Lec.
 solitarius Say.
 brevilabris Lec.
 var. *C. consimilis* Lec.
 impunctifrons Say.
C. emarginatus † Kirby.

Anomoglossa Chaud.

- emarginata Chaud. Bull. Mosc.
Chlænienus emarginatus Dej.

Pasimachus Bon.

- validus Lec.
P. punctulatus † Lec.
 elongatus Lec.
 obsoletus Lec.
 costifer Lec. Proc. Acad. Nat. Sc. VII, 79.

Scarites Fabr.

- subterraneus Fabr. cum var.

Clivina Latr.

- bipustulata Dej.
Scarites bipustulatus Fabr.
Scarites quadrimaculatus Beauv.
 postica Lec.

Aspidoglossa Putz.

- subangulata Lec.
Clivina crenata † Dej.
Dyschirius subangulatus Chaud.
Dyschirius humeralis Chaud.
Aspidoglossa fraterna Putz.
Aspidoglossa vicina Putz.
Clivina bipustulata † Say.

Dyschirius Bon.

- sulcatus Lec.
apicalis † Lec.
 sphaericollis Putz.
Clivina sphaericollis Say.

Tachys Knoch.

- vivax Lec.
 var. *T. mendax* Lec.

incurvus Lec.

Bembidium incurvum Say.

var. *Tachys anceps* Lec.

pulchellus Férè.

dolosus Lec.

sequax Lec.

corrusens Lec.

inornatus Lec.

Bembidium inornatum Say.

Tachyta picipes Kirby.

flavicauda Lec.

Bembidium flavicaudum Say.

Bembidium Latr. (emend. Lec.)

punctato-striatum Say.

stigmaticum Dej.

? *sigillare* Say.

inaequale Say.

arcuarium Dej.

laevigatum Say.

Hydrium laevigatum Lec.

coxendix Say.

Odontium coxendix Lec.

americanus Dej.

cordatum Lec.

dorsale Say.

umbratum Lec.

viridicollis Lec.

Notaphus viridicollis Férè.

patruce Dej.

rapidum Lec.

timidum Lec.

pictum Lec.

quadrinaculatum Gyll.

Carabus quadrinaculatus Linn.

Bembidium oppositum Say.

affine Say, Dej.

Bembidium fallax Dej.

bimaculatum Lec.

Peryphus bimaculatus Kirby.

perspicuum Lec.

cautum Lec.

nitidum Lec.

Eudromus nitidus Kirby.

Cychrus Fabr.

elevatus Fabr.

heros Harris.

Nomaretus Lec.

cavieollis Lec., ante, 3.

Carabus Linn.

serratus Say.

C. lineatopunctatus Dej.

Calosoma Fabr.

luxatum Say.

striatulum Lec., ante, 4.

scrutator Fabr.

calidum Fabr.

var.? *C. lepidum* Lec.

obsoletum Say.

C. luxatum † Dej.

triste Lec.

externum Lec.

Carabus externus Say.

Calosoma longipenne Dej.

Elaphrus Fabr.

Clairvillei Kirby.

intermedius Kirby.

californiens Mann.

var.? *E. similis* Lec.

Omophron Latr.

americanum Dej.

O. Sayi Kirby.

nitidum Lec.

DYTISCIDÆ.

Halipius Latr.

fasciatus Aubé.

innuaculicollis Harris.

H. americanus Aubé.

H. impressus † Kirby, teste White, Brit. Mus.

Cat. 3.

Hydroporus Clairv.

punctatus Aubé.

Laccophilus punctatus Say.

cuspidatus Germ.

Hyphidrus notatus Say.

lacustris Say.

H. pulicarius Aubé.

mixtus Lec. Proc. Acad. Nat. Sc. VII, 296.

semirufus Lec. *ibid.*

vittatus Lec. *ibid.*

catascopium Say.

H. interruptus Say.

H. parallelus Say.

concinus Lec. Proc. Acad. Nat. Sc. VII, 297.

patruelis Lec. *ibid.* VII, 298.

nubilus Lec. *ibid.* VII, 298.

discoideus Lec. *ibid.* VII, 299.

Laccophilus Leach.

maculosus Say.

americanus Aubé.

Coptotomus Say.longulus *Lec.***Copelatus** Er.glyphicus *Lec.**Colymbetes glyphicus* Say.*Copelatus 10-striatus* Aubé.**Agabus** Leach.clavatus *Lec.*, ante, 4.griseipennis *Lec.*, ante, 5.obliteratus *Lec.*, ante, 5.spilotus *Lec.*, ante, 5.teniolatus *Lec.**Colymbetes teniolatus* Harris.*Agabus teniatus* Aubé.**Ilybius** Er.laramacus *Lec.*, ante, 4.**Colymbetes** Clairv.binotatus *Harris.**C. maculicollis* Aubé.**Acilius** Leach.ornaticollis *Aubé.**Thermonectes irroratus* Mels.**Eunectes** Er.sticticus *Er.**Dytiscus sticticus* Linn.**Cybister** Curtis.fimbriolatus *Lec.**Dytiscus fimbriolatus* Say.*Cybister dissimilis* Aubé.**Dytiscus** Linn.anxius *Mann.**D. marginicollis* Lec.*Harrisii* Kirby.**GYRINIDÆ.****Dineutes** McLeay.

two species.

HYDROPHILIDÆ.**Helophorus** Fabr.linearis *Lec.* Proc. Acad. Nat. Sc. VII, 357.lineatus *Say.***Laccobius** Er.agilis *Lec.* Proc. Acad. Nat. Sc. VII, 363.*Hydrophilus agilis* Randall.*Laccobius punctulatus* Mels.**Berosus** Leach.fraternus *Lec.* Proc. Acad. Nat. Sc. VII, 364.**Hydrophilus** Geoffr.triangularis *Say.**Hydrophilus lugubris* Motsch.*Stethoxus subsulcatus* Lec. Proc. Acad. Nat.

Sc. VII, 221.

lateralis *Fabr.**H. nimbatus* Say.sublævis *Lec.* Proc. Acad. Nat. Sc. VII, 363.glaber *Herbst.***Philhydrus** Solier.nebulosus *Lec.**Hydrophilus nebulosus* Say.diffusus *Lec.* Proc. Acad. Nat. Sc. VII, 371.perplexus *Lec.* *ibid.*cinctus *Lec.**Hydrophilus cinctus* Say.*Philhydrus limbatus* Mels.**Hydrobius** Leach.subcupreus *Lec.**Hydrophilus subcupreus* Say.**STAPHYLINIDÆ.¹****Falagria** Mann.venustula *Er.***Xantholinus** Dahl.obscurus *Er.***Staphylinus** Linn.villosus *Grav.*cinnamopterus *Grav.*

¹ Besides the described species of this family mentioned above, nondescripts of the following genera are contained in my collection, but the description of them at present would be attended with no advantage to science, on account of the multitude of other species that remain unknown: they must therefore await the monographing of the entire group.

Myrmedonia, Homalota, Tachyusa, Conurus, Tachinus, Boletobius, Philonthus, Acylophorus, Quediis, Lathrobium, Stenus, Bledius, Oxytelus, Trogophloeus, Apocellus, Boreaphilus, Omalinus.

Philonthus Leach.hepaticus *Er.***Acylophorus** Nordm.flavicollis *Sachse.***Sunius** Leach.longiusculus *Er.**Pæderus longiusculus* Mann.*Pæderus discopunctatus* Say.? binotatus *Say* (*Pæderus*), (fide *Say*).**Pæderus** Fabr.littorarius *Grav.***Lithocharis** Er.confluens *Er.**Lathrobium confluentum* Say.**Stenus** Latr.egenus *Er.*flavicornis *Er.*punctatus *Er.***Euæsthetus** Grav.americanus *Er.***Bledius** Leach.pallipennis *Er.* (fide *Say*).*Orytelus pallipennis* Say.armatus *Er.* (fide *Say*).*Orytelus armatus* Say.melanocephalus *Er.* (fide *Say*).*Orytelus melanocephalus* Say.fasciatus *Er.* (fide *Say*).*Orytelus fasciatus* Say.**Osorius** Leach.latipes *Er.***Anthophagus** Grav.brunnens *Say.***Glyptoma** Er.costale *Er.***PSELAPHIDÆ.****Tyrus** Aubé.humeralis *Lec.**Hamotus humeralis* Aubé, teste Schaum.*Tyrus compar* Lec.**Bryaxis** Leach.rubicunda *Aubé.***SILPHIDÆ.****Necrophorus** Fabr.mediatus *Fabr.*marginatus *Fabr.*Melsheimeri *Kirby.*pustulatus *Herschel.**N. bicolor* Newman.orbicollis *Say.**N. Hallii* Kirby.velutinus *Fabr.**N. tomentosus* Weber.**Silpha** Linn.surinamensis *Fabr.*lapponica *Herbst.**Silpha caudata* Say.*Silpha tuberculata* Germ.*Silpha californica* Mann.*Oicoptoma granigera* Chev.truncata *Say.*peltata *Lec. Proc. Acad. Nat. Sc. VI, 279.**Scarabæus peltatus* Catesby.*Silpha americana* Linn.ramosa *Say.*bituberosa *Lec.*, ante, 6.**Catops** Fabr.simplex *Say* (fide *Say*).basillaris *Say* (fide *Say*).**Agathidium** Illiger.exiguum *Mels.**A. ruficornis* Lec.? pallidum *Say* (fide *Say*).**PHALACRIDÆ.****Phalacrus** Payk.penicillatus *Say.*seriatus *Lec. Proc. Acad. Nat. Sc. VIII, 15.*simplex *Lec. ibid.***Olibrus** Er.pallipes *Lec. Proc. Acad. Nat. Sc. VIII, 17.**Phalacrus pallipes* Say.striatulus *Lec. Proc. Acad. Nat. Sc. VIII, 16.*semistriatus *Lec. ibid.***NITIDULIDÆ.****Carpophilus** Leach.caudalis *Lec. Proc. Acad. Nat. Sc. 1859, 70.*apicalis *Lec. ante, 6.*

carbonatus *Lec.*, ante, 6.

pallipennis *Lec.*

Cercus pallipennis Say.

Carpophilus floralis Er.

Epuræa Er.

rufa *Er.*

Nitidula rufa Say.

Nitidula Fabr. (emend. Er.)

ziezac *Say.*

uniguttata *Mels.*

Omosita Er.

colon *Er.*

Nitidula colon Fabr.

Phenolia Er.

grossa *Er.*

Nitidula grossa Fabr.

Meligethes Steph.

ruficornis *Lec.*, ante, 6.

sævus *Lec.*, ante, 6.

Pocadius Er.

helvolus *Er.*

Pallodes Er.

silaceus *Er.*

Cryptarcha Er.

strigata *Er.*

Nitidula strigata Fabr.

Ips Fabr.

sanguinolentus *Oliv.*

quadrisingnatus *Say.*

bipunctatus *Say.*

Colydium bipunctatum Say.

Trogosita Oliv.

castanea *Mels.*

corticalis *Mels.*

LATHRIDIIDÆ.

Lathridius Herbst.

? 8-dentatus *Say* (fide *Say*).

? *Corticaria denticulata* Kirby.

COLYDIIDÆ.

Bothrideres Er.

geminatus *Er.* (nec *Hald.*)

Lyctus geminatus Say.

Cerylon Latr.

castaneum *Say.*

unicolor *Lec.*

Lathridius unicolor Ziegler.

CUCUJIDÆ.

Læmophleus Er.

biguttatus *Mels.*

Cucujus biguttatus Say.

Læmophleus bisignatus Guér.

Brontes Fabr.

dubius *Fabr.*

Silvanus Latr.

planatus *Germ.*

dentatus *Say.*

Lyctus dentatus Fabr.

DERMESTIDÆ.

Dermestes Linn.

marmoratus *Say.*

nubilus *Say.*

caninus *Germ.*

clongatus *Lec.*

vulpinus *Fabr.*

HETEROCERIDÆ.

Heterocerus Fabr.

pallidus *Say* (fide *Say*).

pusillus *Say* (fide *Say*).

GEORYSSIDÆ.

Georyssus Latr.

pusillus *Lec.* Proc. Acad. Nat. Sc. VI, 44.

SCAPHIDIIDÆ.

Scaphidium Oliv.

4-pustulatum *Say.*

HISTERIDÆ.

Hololepta Payk.

fossularis *Say.*

æqualis *Say* (fem.)

lucida *Lec.*

Hister Linn.

- instratus *Lec.*, ante, 7.
 biplagiatus *Lec.*
 dispar *Lec.*
 deparator *Say.*
 abbreviatus *Fabr.*
 nubilus *Lec.*, ante, 7.
 pollutus *Lec.*, ante, 7.
 americanus *Payk.*
 subrotundus *Er.*
 carolinus *Payk.*
 Lecoutei *Lec.*
Platysoma depressum Er., *Lec.*
Platysoma Lecoutei Marseul.
 parallelus *Say.*

Paromalus Er.

- aequalis *Er.*
Hister aequalis *Say.*
 bistriatus *Er.*

aprinus Leach.

- lugens *Er.*
californicus *Mann.*
 spureus *Lec.*, ante, 7.
 parumpunctatus *Lec.*, ante, 7.
 pratensis *Lec.*, ante, 8.
 pennsylvanicus *Er.*
Hister pennsylvanicus *Payk.*
 patruelis *Lec.*

Plegaderus Leach.

- transversus *Lec.*
Hister transversus *Say.*

Acritus Lec.

- exiguus *Lec.*
Abreus criguus *Er.*

SCARABÆIDÆ.**Xyloryctes** Hope.

- satyrus *Burm.*
Geotrupes satyrus *Fabr.*

Phileurus Latr.

- valgus *Drj.*
Geotrupes valgus *Fabr.*
Phileurus castaneus *Hald.*

Strategus Hope.

- mormon *Burm.*

Aphonus Lec.

- pyriformis *Lec.* *Proc. Acad. Nat. Sc. VIII, 21.*
Bolhynus pyriformis *Lec.*

tridentatus *Lec.* *ibid.* VIII, 22.

- Scarabæus tridentatus* *Say.*
Bolhynus tridentatus *Lec.*

Ligyris Burm.

- gibbosus *Lec.* *Proc. Acad. Nat. Sc. VIII, 20.*
Scarabæus gibbosus *DeGeer.*
Podalgus variolosus *Burm.*
 relictus *Lec.* *Proc. Acad. Nat. Sc. VIII, 21.*
Scarabæus relictus *Say.*
Heteronychus relictus *Burm.*

Osmoderma Lepell.

- cremicola *Gory.*
Cetonia cremicola *Knoch.*

Cremastochilus Knoch.

- nitens *Lec.* *Proc. Acad. Nat. Sc. VI, 232.*
 Knoehii *Lec.* *ibid.* VI, 231.
 saucius *Lec.* *Journ. Acad. Nat. Sc. 2d ser. IV, 16.*

Euryomia Burm. (emend. Lac.)

- inda *Lac.* *Gen. Col. III, 528.*
Scarabæus indus *Linn.*
Cetonia barbata *Say.*
Erirhipis inda *Burm.*
 melancholica *Lac.* *Gen. Col. III, 527.*
Cetonia melancholica *Gory.*
Euphoria melancholica *Schaum.*
 sepulchralis *Lac.* *ibid.*
Cetonia sepulchralis *Fabr.*
Cetonia Reichii *Gory.*
Euphoria sepulchralis *Burm.*
 fulgida *Lac.* *Gen. Col. III, 528.*
Cetonia fulgida *Fabr.*
Erirhipis fulgida *Fabr.*
 areata *Lac.* *Gen. Col. III, 528.*
Cetonia areata *Fabr.*
Stephanucha areata *Burm.*

Allorhina Burm. (emend. Lac.)

- nitida *Lac.* *Gen. Col. III, 497.*
Scarabæus nitidus *Linn.*
Cotinis nitida *Burm.*

Anomala Samouelle (emend. Burm.).

- minuta *Burm.*
 marginata *Burm.*
Melolontha marginata *Fabr.*
Melolontha annulata *Germ.*
 varians *Burm.*
Melolontha varians *Fabr.*

Strigoderma Burm.

arboricola Burm.

Melolontha arboricola Fabr.**Polyphylla** Harris.

decemlineata Lec. (infra).

Melolontha 10-lineata Say.

Hammondi Lec. J. Ac. Nat. Sc. 2d ser. III, 228.

Lachnosterna Hope.

lanceolata Lec. J. Ac. Nat. Sc. 2d ser. III, 237.

Melolontha lanceolata Say.*Postcogoptera lanceolata* Blanchard.

frontalis Lec. J. Ac. Nat. Sc. 2d ser. III, 239.

longitarsis Lec. ibid.

Melolontha longitarsis Say.

futilis Lec.

fusca Lec. Journ. Acad. Nat. Sc. 2d ser. III, 244.

Melolontha fusca Fröhl.*Melolontha quercina* Knoch.*Melolontha fervens* Gyll.*Melolontha fervida* Oliv.

cephalica Lec. J. Ac. Nat. Sc. 2d ser. III, 245.

fraterna Lec. ibid. III, 249.

Phyllophaga fraterna Harris.*Ancylonycha fraterna* Blanch.

rugosa Lec. ibid. III, 252.

Ancylonycha rugosa Mels.

affinis Lec. ibid. III, 252.

ciliata Lec. ibid. III, 253.

hirticula Hope.

Melolontha hirticula Knoch.*Melolontha hirsuta* Say.

robusta Lec. ibid. III, 257.

crenulata Lec. ibid. III, 258.

Melolontha crenulata Fröhl.*Melolontha georgicana* Gyll.*Phyllophaga georgicana* Harris.*Ancylonycha crenulata* Blanch.

glabricula Lec. ibid. III, 260.

tristis Lec. ibid. III, 261.

Melolontha tristis Fabr.*Melolontha pilosicollis* Knoch.*Trichesthes pilosicollis* Er.*Trichestes tristis* Burm.**Listrochelus** Blanch.

obtusus Lec. Journ. Acad. Nat. Sc. 2d ser. III, 264.

falsus Lec. ibid.

fimbripes Lec. ibid.

Diplotaxis Kirby.

obscura Lec., ante, 9.

Harperi Blanch.

frondicola Lec.

Melolontha frondicola Say.*Diplotaxis testacea* Burm.

truncatula Lec. J. Ac. Nat. Sc. 2d ser. III, 269.

morula Lec. ibid. III, 270.

Haydenii Lec. ibid. III, 272.

innocia Lec. ibid. III, 273.

Diazus Lec.

rudis Lec., ante, 10.

Dichelonycha Kirby.

truncata Lec. J. Ac. Nat. Sc. 2d ser. III, 281.

Serica M'Leay.

vespertina Dej.

Melolontha vespertina Gyll. Say.*Camptorhina atricapilla* Kirby.

sericea Burm.

Melolontha sericea Say.

curvata Lec. J. Ac. Nat. Sc. 2d ser. III, 276.

Macroductylus Latr.

angustatus Lec. J. Ac. Nat. Sc. 2d ser. III, 278.

Melolontha elongata Herbst.*Melolontha angustata* Beauv.*Macroductylus polyphagus* Burm.**Hoplia** Illiger.

laticollis Lec. J. Ac. Nat. Sc. 2d ser. III, 284.

Geotrupes Latr.

opacus Hald. Proc. Acad. Nat. Sc. 2d ser. VI, 362.

Odontæus Klug.

flicicornis Say.

Geotrupes flicicornis Say.**Bolbocerus** Kirby.

lazarus Klug.

Geotrupes lazarus Fabr.*Geotrupes melibæus* Fabr.**Canthon** Hoffmans.

levis Mels.

Scarabæus pilularius De Geer.*Scarabæus lævis* Drury.*Ateuchus volvens* Fabr.

chalcites Mels.

Coprobis chalcites Hald.

ebenus Mels.

Ateuchus ebenus Say.

nigricornis Mels.

Ateuchus nigricornis Say.

praticola Lec., ante, 10.

viridis Mels.

Ateuchus viridis Beauv.

Onthophagus viridicatus Say.

Ateuchus obsoletus Say.

Onthophagus Latr.

orpheus.

Scarabæus orpheus Panzer.

Copris canadensis Fabr.

hecate.

Scarabæus hecate Panzer.

Copris latebrosus Fabr.

Copris hastator Fabr.

Copris oblectus Beauv.

Phanæus M'Leay.

carnifex M'Leay.

Scarabæus carnifex Linn.

triangularis Lec.

Copris triangularis Say.

Phanæus torrens Lec.

Copris Geoffr.

anaglypticus Say.

ammon Fabr.

Ochodæus Latr.

musculus Lec.

Bolbocerus musculus Say.

Ochodæus americanus Westwood.

Aphodius Illiger.

denticulatus Hald.

curtus Hald.

granarius Illiger.

Scarabæus granarius Linn.

Aphodius 4-tuberculatus Fabr.

vittatus Say.

femoralis Say.

concavus Say.

A. lævigatus Hald.

oblongus Say.

A. badipes Mels.

Euparia Lep. (emend. Er.)

stercorator Er.

Aphodius stercorator Fabr.

strigata Lec.

Aphodius strigatus Say.

Trox Fabr.

alternans Lec. Proc. Acad. Nat. Sc. VII, 211.

tuberculatus Ol.

Scarabæus tuberculatus DeG.

Trox canaliculatus Say.

Trox serrulatus Beauv.

sordidus Lec. ibid VII, 211.

capillaris Say.

atrox Lec. Proc. Acad. Nat. Sc. VII, 214.

Omorgus Er.

scutellaris Lec. Proc. Acad. Nat. Sc. VII, 214.

Trox scutellaris Say.

pustulatus Lec.

Trox tuberculatus† Beauv.

punctatus Lec. Proc. Acad. Nat. Sc. VII, 215.

Trox punctatus Germ.

Trox alternatus Say.

erinaceus Lec.

morsus Lec. ibid. VII, 216.

Lucanus Linn.

capreolus Linn.

L. dama Thunberg.

placidus Say.

L. lentus Lap. Hist. Nat. II, 171.

Doreus M'Leay.

parallelus Burm.

Lucanus parallelus Say.

Platycerus Geoffr.

quercus Schönh.

Lucanus quercus Weber.

Platycerus securidens Say.

BUPRESTIDÆ.

Ancylochira Esch.

confluens Lec.

Buprestis confluenta Say.

mauliventris Lec.

Buprestis maculiventris Say.

Buprestis serotata Lap.

Anoplis rusticorum Kirby.

Chalcophora Sol.

eampestris Lec. (fide Say).

Buprestis campestris Say.

Buprestis substrigosa Lap. & Gory.

Melanophila Esch.

longipes Lec.

Apatura appendiculata† Lap. & Gory.

Bup. (Oxypteris) appendiculata† Kirby.

Melanophila immaculata Gory.

atropurpurea Lec.

Buprestis atropurpurea Say.

fulvoguttata *Lec.*

- Buprestis fulvoguttata* Harris.
Apatura octospilota Lap. & Gory.
Apatura croceosignata Lap. & Gory.
Apatura decolorata Lap. & Gory.

Anthaxia Esch.

- quercata *Dej.*
Buprestis quercata Fabr.
 viridicornis *Dej.*
Buprestis viridicornis Say.

Chrysobothris Esch.

- sexguttata *Lec.*
Buprestis sexguttata Say.

Psiloptera Sol.

- Woodhousei *Lec.*
Dicerca? *Woodhousei* Lec. Proc. Acad. Nat.
 Sc. VI, 68.
 var. *P. valens* Lec. *ibid.* 1858, 66.

Dicerca Esch.

prolongata *Lec.*

Poeilonota Esch.

- cyanipes *Lec.*
Buprestis cyanipes Say.

Acmæodera Esch.

mixta *Lec.* Trans. Am. Phil. Soc. XI, 227.

Ptosima Serv.

- gibbicollis *Lec.*
Buprestis gibbicollis Say.
Ptosima luctuosa Gory.

Agrilus Lap.

- bilineatus *Say.*
Buprestis bilineata Weber.
Agrilus bivittatus Kirby.
Agrilus flavolineatus Mann.
Agrilus aurolineatus Gory.
 latebrus *Lap.*
 granulatus *Say.*
 politus *Say.*
 defectus *Lec.* Trans. Am. Phil. Soc. XI, 244.
 pusillus *Say.*
 obolinus *Lec.* Trans. Am. Phil. Soc. XI, 248.
 otiosus *Say.*
 lateralis *Say.*

Brachys Sol.

- terminans *Lap.*
Buprestis terminans Fabr.

ELATERIDÆ.

Tharops Lap.

- ruficornis *Lec.* Trans. Am. Phil. Soc. X, 411.
Melasis ruficornis Say.

Hylochares Latr.

- nigricornis *Lec.* Trans. Am. Phil. Soc. X, 413.
Melasis nigricornis Say.

Microrhagus Esch.

- triangularis *Lec.* Trans. Am. Phil. Soc. X, 419.
Elater triangularis Say.

Pedetes Kirby.

- eucllatus *Lec.* Trans. Am. Phil. Soc. X, 425.
Elater eucllatus Say.
Athous hypoleucus Mels.
Athous procericollis Mels.
Athous strigatus Mels.

Limonius Esch.

- auripilis *Lec.* Trans. Am. Phil. Soc. X, 429.
Elater auripilis Say.
 quercinus *Dej.*
Elater quercinus Say.
 basillaris *Lec.* Trans. Am. Phil. Soc. X, 431.
Elater basillaris Say.

Cratonychus Er.

- macer *Lec.* Trans. Am. Phil. Soc. X, 473.
 incertus *Lec.* Trans. Am. Phil. Soc. X, 474.
 clandestinus *Er.*
 fissilis *Lec.*
Cratonychus laticollis Er.
 ?*Elater brevicollis* Herbst.
Elater cinereus (fissilis) Say.
Elater (Melanotus) cinereus Harris.
Cratonychus ochraceipennis Mels.
Cratonychus sphenoidalis Mels.
 communis *Er.*
Elater communis Gyll.
Elater cinereus Weber.
 cribulosus *Lec.* Trans. Am. Phil. Soc. X, 478.

Monocrepidius Esch.

- vespertinus *Dej.*
Elater vespertinus Fabr.
Elater finitimus Say.
Monocrepidius serotinus Germ.
 auritus *Germ.*
Elater auritus Herbst.
Oophorus crassicollis Mels
 bellus *Dej.*
Elater bellus Say.
Cryptohypnus bellus Germ.

Æolus Esch.

- dorsalis *Candèze*, Mon. Elat. II, 285.
Elater dorsalis Say.
Cryptohypnus dorsalis Germ.

Ædostethus Lec.

- femoralis *Lec.* Trans. Am. Phil. Soc. X, 489.

Adlocera Latr.

- impressicollis *Lec.* Trans. Am. Phil. Soc. X, 490.
Elater impressicollis Say.
Elater lepturus Say.
Adlocera senilis Germ.
marmorata *Germ.*
Elater marmoratus Fabr.

Lacon Lap.

- rectangularis *Candèze*, Mon. El. I, 155.
Elater rectangularis Say.

Melanactes Lec.

- piceus *Lec.* Trans. Am. Phil. Soc. X, 494.
Elater piceus DeGeer.
Elater lævigatus Fabr.
Elater morio (var.) Say.
Pristilophus lævigatus Germ.
Pristilophus femoralis Mels.
puncticollis *Lec.* Trans. Am. Phil. Soc. X, 495.
Pristilophus puncticollis *Lec.* Proc. Ac. Nat.
 Sc. VI, 68.

Alaus Esch.

- oculatus *Esch.*
Elater oculatus Fabr.
myops *Esch.*
Elater myops Fabr.
Elater luscus Oliv.
gorgops *Lec.* Journ. Acad. Nat. Sc. 2d ser. IV, 35.
Alaus oculatus (var.) *Lec.* Trans. Am. Phil.
 Soc. 10, 496.

Cardiophorus Esch.

- erythropus *Er.*
Cardiophorus amictus Mels.
 ? *Elater convexus* Say.

CYPHONIDÆ.**Helodes** Latr.

- ruficollis *Lec.*
Lampyrus ruficollis Say.
Elodes fragilis Ziegler.

Scirtes Illiger.

- centralis *Lec.*
Altica centralis Say.
 ? *Scirtes orbiculatus* Fabr.
Scirtes suturalis Ziegler.
Scirtes lateralis *Lec.*

LYCIDÆ.**Calopteron** Lap.

- typicum *Lec.*
Digrapha typica Newm.
Digrapha discrepans Newm.
Digrapha affinis *Lec.*
terminale *Lec.*
Lycus terminalis Say.

Temnostoma Guér.

- sanguinipennis *Lec.*
Lycus sanguinipennis Say.

LAMPYRIDÆ.**Photinus** Lap. (emend. Lac.)§ **Ellychnia** Lec.

- nigricans *Lec.*
Lampyrus nigricans Say.
 ? *Lampyrus obscura* Fabr.
corrusca *Lec.*
Lampyrus corrusca Linn.

§§ **Photinus** Lap.

- pyralis *Lap.*
Lampyrus pyralis Linn.
Lampyrus centrata Say.
Lampyrus rosata Germ.
marginella *Lec.*
punctulata *Lec.*
obseurella *Lec.*

Photuris Lec.

- pennsylvanica *Lec.*
Lampyrus pennsylvanica DeGeer.
Lampyrus versicolor Fabr.
divisa *Lec.*

TELEPHORIDÆ.**Chauliognathus** Hentz.

- marginatus *Hentz.*
Cantharis marginata Fabr.
Cantharis ligata Say.
basalis *Lec.*, ante, 13.

Telephorus Geoffr.

- collaris *Lec.*
 bilineatus *Lec.*
 Cantharis bilineatus Say.
 jaectatus *Lec.*
 Cantharis jaectata Say.
 carolinus.
 Cantharis carolina Fabr.
 luteicollis *Germ.*
 T. cinctellus Lec.
 dichrous *Lec.*
 flavipes *Lec.*

Podabrus Fischer.

- rugosulus *Lec.*
 punctulatus *Lec.*

Trypherus Lec.

- latipennis *Lec.*
 Malthinus latipennis Germ.
 Molorchus marginalis Say.
 Malthinus marginalis Say.
 Lygerus latipennis Kiesenwetter.

MELYRIDÆ.**Collops** Er.

- bipunctatus *Er.*
 Malachius bipunctatus Say.
 tricolor *Er.*
 Malachius tricolor Say.
 punctatus *Lec.* Proc. Acad. Nat. Sc. VI, 164.
 quadrimaculatus *Er.*
 Malachius 4-maculatus Fabr.
 confluent *Lec.* Proc. Acad. Nat. Sc. VI, 164.
 punctulatus *Lec.* *ibid.*

Ebæus Er.

- morulus *Lec.* Proc. Acad. Nat. Sc. VI, 167.

Dasytes Fabr.

- senilis *Lec.* Proc. Acad. Nat. Sc. VI, 170.

Byturus Fabr.

- unicolor *Say.*

CLERIDÆ.**Cymatodera** Gray.

- undulata *Lec.*
 Tillus undulatus Say.
Cymatodera longicollis Spin.
Cymatodera Bosei Chevr.

Trichodes Herbst.

- ornatus *Say.*
 Nuttalli *Kirby.*

Clerus Geoffr.

- analis *Say.*
 cordifer *Lec.*
 sphegeus *Fabr.*

Hydnocera Newm.

- humeralis *Newman.*
 Clerus humeralis Germ.

Enoplium Latr.

- pilosum *Latr.*
 Lampyrus pilosa Forster.
 var. *Enoplium onustum* Say
 Enoplium marginatum Say.
 quadripunctatum *Say.*

Orthopleura Spin.

- damicornis *Spin.*
 Tillus damicornis Fabr.
 Enoplium thoracicum Say.

Corynetes Fabr.

- rufipes *Fabr.*
 violaceus *Fabr.*

PTINIDÆ.**Dorcatoma** Herbst.

- simile *Say.*

BOSTRICHIDÆ**Bostrichus** Geoffr. (emend. Læe.)

- bicaudatus *Lec.*
 Apate bicaudatus Say.

TENEBRIONIDÆ.**Epitragus** Latr.

- canaliculatus *Say.*

Edrotes Lec.

- rotundus *Lec.*
 Pimelia rotunda Say.

Trimytis Lec.

- pruinosa *Lec.*

Pelecyphorus Sol.

- sordidus *Lec.* Proc. Acad. Nat. Sc. VI, 446.

Euschides Lec.convexa *Lec.*, ante, 14.polita *Lec.**Asida polita* Say.**Asida** Latr.opaca *Say.***Pactostoma** Lec.anastomosis *Lec.* J. Ac. Nat. Sc. 2d ser. IV, 19.*Asida anastomosis* Say.*Ologlyptus anastomosis* Lac.**Eleodes** Esch.obscura *Lec.**Blaps obscura* Say.? *Blaps hispilabris* Say.dispersa *Lec.* Proc. Acad. Nat. Sc. 1858, 182.acuta *Lec.**Blaps acuta* Say.suturalis *Lec.**Blaps suturalis* Say.tricostata *Lec.**Blaps tricostata* Say.*Pimelia alternata* Kirby.sulcata *Lec.* Proc. Acad. Nat. Sc. VI, 67.obsoleta *Lec.**Blaps obsoleta* Say.fusiformis *Lec.* Proc. Acad. Nat. Sc. 1858, 184.extricata *Lec.**Blaps extricata* Say.carbonaria *Lec.**Blaps carbonaria* Say.nigrina *Lec.* Proc. Acad. Nat. Sc. 1858, 186.Haydenii *Lec.* *ibid.* 1858, 186.viator *Lec.* *ibid.* 1858, 188.**Promus** Lec.opacus *Lec.* Say's Ent. Writings, II, 155.*Blaps opaca* Say.**Embaphion** Say.municatum *Say* (Akis).contusum *Lec.* Journ. Acad. Nat. Sc. 2d ser. IV, 40.**Coniontis** Esch.obesa *Lec.***Eusattus** Lec.reticulatus *Lec.**Zophosis reticulata* Say.convexus *Lec.***Pedinus** Latr.? suturalis *Say* (fide *Say*).**Blapstinus** Waterhouse.interruptus *Lec.**Opatrum interruptum* Say.*Blapstinus æneolus* Mels.pratensis *Lec.*, ante, 15.vestitus *Lec.*, ante, 15.**Centronopus** Sol.opacus *Lec.*, ante, 15.**Upis** Fabr.levis *Oliv.***Nyctobates** Esch.pennsylvanicus *Lec.**Tenebrio pennsylvanicus* DeGeer.*Upis chrysops* Herbst.barbatus *Lec.**Upis barbatus* Knoch.**Boletophagus** Illiger.? cornutus *Fabr.**Opatrum cornutum* Panz.*Opatrum bifurcum* Fabr.**Uloma** Redt.culinaris *Redt.**Tenebrio culinaris* Linn.**Tenebrio** Linn.tenebrioides *Lec.**Helops tenebrioides* Beauv.*Tenebrio badipes* Mels.**Paratenetus** Spin.punctatus *Spin.***Adelina** Dej.pallida *Lec.**Pytho pallida* Say.**Diaperis** Geoffr.hydri *Fabr.***Platydema** Lap.excavatum *Lap.***Allecula** Fabr.punctulata *Mels.*obscura *Say.***MELANDRYADÆ.****Eustrophus** Fabr.bicolor *Fabr.*

Melandrya Fabr.labiata *Say.***PEDILIDÆ.****Scraptia** Latr.plagiata *Mels.**S. americana* Hald.**Stereopalpus** Ferté.guttatus *Lec. Proc. Acad. Nat. Sc. VII, 271.***ANTHICIDÆ.****Notoxus** Geoffr.anchora *Hentz.*serratus *Lec.*monodon *Ferté.**Anthicus monodon* Fabr.marginatus *Lec.*subtilis *Lec.***Anthicus** Payk.elegans *Ferté.*rejeetus *Lec.*cervinus *Ferté.**A. bifasciatus* || *Say.**A. terminalis* *Lec.**A. bizonatus* *Ferté.***MORDELLIDÆ.****Mordella** Fabr.quadripunctata *Lec.**Anaspis 4-punctata* *Say.*insulata *Lec.*, ante, 16.marginata *Mels.***Mordellistena** Costa.æmula *Lec.*, ante, 16.divisa *Lec.*, ante, 17.**RHIPIPHORIDÆ.****Emenadia** Lap.Sayi *Lec.**Rhipiphorus bicolor* || *Say.*limbatus *Lac.**Rhipiphorus limbatus* Fabr.pectinatus *Lec.**Rhipiphorus pectinatus* Fabr.*Rhipiphorus varicolor* Gerst.**Myodites** Latr.scaber *Lec. Proc. Acad. Nat. Sc. VI, 67.***MELOIDÆ.****Henous** Hald.confertus *Lec. Proc. Acad. Nat. Sc. VI, 330.**Meloe conferta* *Say.**Henous techanus* Hald.**Lytta** Fabr.reticulata *Say.*Nuttalli *Say.*var. *Cantharis fulgifer* *Lec.*sphaericollis *Say.*Engelmanni *Lec. Proc. Acad. Nat. Sc. VI, 337.**Pyrota Engelmanni* *Lec.*discoidea *Lec. Proc. Acad. Nat. Sc. VI, 338.*pennsylvanica *Lec. ibid. VI, 339.**Cantharis pennsylvanica* DeGeer.*Lytta atrata* Fabr.ferruginea *Say.*maculata *Say.*conspersa *Lec. Proc. Acad. Nat. Sc. VI, 340.*segmentata *Say (segmenta).*albida *Say.**L. luteicornis* *Lec.*immaculata *Say.*♂ *L. articularis* *Say.*longicollis *Lec. Proc. Acad. Nat. Sc. VI, 343.*Fabricii *Lec. ibid.**L. cinerica* || Fabr.**Nemognatha** Illiger.bicolor *Lec. Proc. Acad. Nat. Sc. VI, 345.*lurida *Lec. ibid.*lutea *Lec. ibid. VI, 346.*piezata *Lec. ibid. VI, 347.**Zonitis piezata* Weber, Fabr.*Zonitis vittata* Fabr.immaculata *Say.*minima *Say.***Zonitis** Fabr.atripennis *Lec. Proc. Acad. Nat. Sc. VI, 349.**Nemognatha atripennis* *Say.***CEDEMERIDÆ.****Asclera** Schmidt.puncticollis *Hald.**Edemera puncticollis* *Say.*

ruficollis *Hald.*

Edemera ruficollis Say.

? vestita *Say* (*Edemera*), fide *Say*.

(an potius *Stereopalpus*.)

CURCULIONIDÆ.

Bruchus Linn.

discoideus *Say*.

and five others.

Spermophagus Steven.

Robiniæ *Sch.*

Bruchus Robiniæ Fabr.

Cratoparis Sch.

lunatus *Sch.*

Anthrribus lunatus Fabr.

Anthrribus marmoratus Ol.

Attelabus Linn.

nigripes *Lee*.

Pterocolus Schönh.

ovatus *Schönh.*

Attelabus ovatus Fabr.

Rhynchites Herbst.

bicolor *Herbst.*

Attelabus bicolor Fabr.

æneus *Boheman.*

æreatus *Say*.

Apion Herbst.

four species.

Ophryastes Schönh.

latirostris *Lee*. Proc. Acad. Nat. Sc. VI, 443.

ligatus *Lee*. *ibid.*

sulcistrostris *Schönh.*

Liparus sulcistrostris Say.

vittatus *Schönh.* (*infra*).

Liparus vittatus Say.

Epicærus Schönh.

? imbricatus (fide *Say*).

Liparus imbricatus Say.

and four species? with four others of allied genera.

Platyomus Schönh.

auriceps *Schönh.*

Curculio auricephalus Say.

Thylacites Germ.

microsus *Schönh.*

T. microps Say.

Tanymecus Germ.

canescens *Schönh.*

confertus *Schönh.*

T. confusus Say.

Cleonus Schönh.

pulvereus *Lee*., ante, 18.

trivittatus *Say*.

angularis *Lee*., ante, 18.

Listroderes Schönh.

three species.

Lepyryus Germ.

geminatus *Say*.

Lithodus Germ.

humeralis *Germ.*

Brachycerus humeralis Say.

Thecsternus humeralis Say.

rectus *Lee*. Proc. Acad. Nat. Sc. VIII, 18.

affinis *Lee*. Proc. Acad. Nat. Sc. VIII, 18.

rudis *Lee*. Proc. Acad. Nat. Sc. VIII, 18.

crossus *Lee*. Proc. Acad. Nat. Sc. VIII, 18.

longior *Lee*. Proc. Acad. Nat. Sc. VIII, 19.

morbillosus *Lee*. Proc. Acad. Nat. Sc. VIII, 19.

Lixus Fabr.

two species.

Eirrhinus Schönh.

and allied genera 15 species.

Grypidius Schöuh.

one species.

Balaninus Germ.

one species.

Piazorhinus Schönh.

scutellaris *Schönh.* Cure. III, 472; VII, 2, 352.

Attelabus scutellaris Say.

Baridius Schönh.

seven species.

Centrinus Schönh.

two species.

Ceuthorhyncus Schüppel.

four species.

Rhyssematus Schönh.

lineaticollis *Schönh.*

Rhynchanus lineaticollis Say.

Acalles Schönh.

three species.

Conotrachelus Latr.

posticatus *Schönh.*

Sphenophorus Schönh.

pulchellus *Schönh.*
cultirostris *Germ.*
compressirostris *Say.* (Calandra).
and seven others.

Cossonus Clairv.

subareatus *Schönh.*
and one other species.

Tomicus Latr.

pini *Harris.*
Bostrichus pini *Say.*
caligraphus *Germ.*
Bostrichus excusus *Say.*

CERAMBYCIDÆ.**Mallophon** Serv.

dasystemus *Hald.*
Prionus dasystemus *Say.*
ciliipes *Hald.*
Prionus ciliipes *Say.*

Prionus Geoffr.

palparis *Say.*
imbriicornis *Oliv.*
fissicornis *Hald.*
integer *Lec.*
emarginatus *Say.*

Criocephalus Muls.

productus *Lec.*
asperatus *Lec.*, ante, 19.
agrestis *Hald.*
Callidium agreste *Kirby.*

Semanotus Muls.

brevilineus *Lec.*
Callidium brevilineum *Say.*
Physocnemum brevilineum *Hald.*

Callidium Fabr.

variabile *Fabr.*
amœnum *Say.*

Dryobius Lec.

sexfasciatus *Lec.*
Callidium serfasciatum *Say.*

Heliomanes Newman.

bimaculatus *Newman.*
Molorchus bimaculatus *Say.*
Molorchus affinis *Lec.*
Heliomanes obscurus *Lec.* (err. typog.)

Eburia Serv.

quadrigeminata *Hald.*
Stenocorus quadrigeminatus *Say.*

Elaphidion Serv.

simplicicollis *Hald.*
E. pulverulentum † *Hald.*
mucronatum *Newman.*
Stenocorus mucronatus *Say.*
villosum *Dej.*
Stenocorus villosus *Fabr.*
Stenocorus putator *Peck.*

Eriphus Serv.

ignicolle *Lec.*
Callidium ignicolle *Say.*
Callidium sanguinicolle *Germ.*

Arhopalus Serv. (emend. *Lec.*)

fulminans *Serv.*
Clytus fulminans *Fabr.*
eharus *Lec.*
Clytus charus *Say.*
pictus *Lec.*
Cerambyx pictus *Drury.*
Leptura Robiniæ *Forster.*
Clytus flexuosus *Fabr.*

Rhopalophorus Serv.

longipes *Lec.*
Stenocorus longipes *Say.*
Tinopus longipes *Lec.*

Purpuricenus Serv.

humeralis *Dej.*
Cerambyx humeralis *Fabr.*

Clytus Fabr.

scutellaris *Dej.*
Callidium scutellare *Oliv.*
erythrocephalus *Fabr.*
undulatus *Say.*
C. Sayi *Lap.*
C. undatus *Kirby.*
capræa *Say.*
C. elevatus *Lap.*

Acmæops Lec.

- bivittata Lec.
Leptura bivittata Say.
 dorsalis Lec., ante, 21.

Typocerus Lec.

- sinuatus Lec.
Leptura sinuata Newman.
Stenura 8-notata Hald.

Leptura Linn.

- cribripennis Lec., ante, 21.
 rubrica Say.

Monilema Say.

- annulatum Say.

Plectrodera Lec.

- scalator Lec.
Lamia scalator Fabr.
Lamia Belli Lec.

Oberea Muls.

- perspicillata Hald.
 oculaticollis Lec.
Saperda oculaticollis Say.

Stenostola Muls.

- pergrata Lec.
Saperda pergrata Say.
 gentilis Lec.
 saturnina Lec., ante, 21.

Amphionycha Lec.

- ardens Lec., ante, 22.

Tetrops Kirby.

- canescens Lec.

Tetraopes Dalman.

- tetraophthalmus Harris.
Cerambyx tetraophthalmus Forster.
Lamia tornator Fabr.
 femoratus Lec.
 annulatus Lec.

Saperda Fabr.

- calcarata Say
 var. *S. adspersa* Lec.
 mutica Say.
 discoidea Fabr.
 puncticollis Say.
S. trigeminata Randall.
 ?inornata Say (fide Say).

Pogonocherus Latr.

- parvulus Lec.

Psenocerus Lec

- supernotatus Lec.
Clytus supernotatus Say.

Leptostylus Lec.

- aculiferus Lec.
Lamia aculifera Say.
Anniscus marginellus Hald.

Liopus Serv.

- cinereus Lec.

Acanthoderes Serv.

- decipiens Lec.
Ergomorphus decipiens Hald.

CHRYSOMELIDÆ.**Clythra** Laichart.

- laticlavia Sch.
Chrysomela laticlavia Forster.
Clythra obsita Fabr.

Babia Lac.

- quadriguttata Lac.
Clythra 4-guttata Oliv.

Coscinoptera Lac.

- franciscana Lec., ante, 22.

Cryptocephalus Geoffr.

- lativittis Germ.
C. geminatus Hald.
 guttulatus Oliv., Suffr.
C. lautus Newman.
 mucoreus Lec., ante, 23.
 notatus Oliv.
C. quadrimaculatus Say, Hald.
 quadriguttulus Suffrian.
 dispersus Hald.
 venustus Fabr., Hald.
C. ornatus Say.
C. calidus Suffr.
 leucomelas Suffrian.
 fasciatus Say.
 amatus Hald.
 confluens Say.
 viridis Hald.
Monachus viridis Mels.
 ?*Cryptocephalus auratus* Fabr.

Pachybrachys Suffr.hepaticus *Hald.**Cryptocephalus hepaticus* Mels.tridens *Mels.*mollis *Hald.*viduatus *Suffr.**Cryptocephalus viduatus* Fabr.*Cryptocephalus bivittatus* Say.

and four? other species.

Colaspis Fabr.favosa *Say.*

and six? other species.

Metachroma Lec.interruptum *Lec.**Colaspis interrupta* Say.pallidum *Lec.**Colaspis pallida* Say.var. *Colaspis dubiosa* Say.**Paria** Lec.sexnotata *Lec.**Colaspis 6-notata* Say.quadrinotata *Lec.**Colaspis 4-notata* Say.aterrima *Lec.**Colaspis aterrima* Oliv.opacicollis *Lec.*, ante, 23.pumila *Lec.*, ante, 23.**Hcteraspis** Lec.nebulosa *Lec.*, ante, 23.smaragdula *Lec.*, ante, 24.**Myochrous** († Chevr.).denticollis *Lec.**Colaspis denticollis* Say.squamosus *Lec.*, ante, 24.**Chrysomela** Linn.sealaris *Lec.*philadelphica *Linn.*multipunctata *Say.*var. *C. verrucosa* Sulfr. Ent. Zeit. 1858, 265.exclamationis *Fabr.*conjuncta *Rogers*, Proc. Acad. Nat. Sc. VIII, 34.disrupta *Rogers*, *ibid.*lunata *Fabr.**C. hybrida* Say.pulehura *Fabr.*var. *C. casta* Rogers, Pr. Ac. Nat. Sc. VIII, 33.*C. lineata* DeGeer.*C. festiva* Fabr.incisa *Rogers*, Proc. Acad. Nat. Sc. VIII, 34.præcelsa *Rogers*, Proc. Acad. Nat. Sc. VIII, 35.auripennis *Say.*basillaris *Say* (fide Say).flavomarginata *Say.*interrupta *Fabr.*scripta *Fabr.*obsoleta *Say.*dissimilis *Say.*formosa *Say.***Doryphora** Fabr.10-lineata *Say.*Rogersii *Lec.* Journ. Ac. Nat. Sc. 2d ser. IV, 26.trimaculata *Say.**Chrysomela trimaculata* Linn.*Chrysomela clivicollis* Fabr.**Blepharida** Rogers († Chevr.).rhois *Rogers.**Chrysomela rhois* Forster.*Chrysomela stolidia* Fabr.*Altica virginica* Fröhlich.**Edionychis** Latr.gibbitarsa *Lec.**A. gibbitarsa* Say.scripticollis *Lec.**A. scripticollis* Say.lobata *Lec.*, ante, 24.**Haltica** Illiger.alternata *Illiger.**A. quinquevittata* Say.punctigera *Lec.*, ante, 24.pluriligata *Lec.*, ante, 25.collaris *Illiger.**Galleruca collaris* Fabr.cervicalis *Lec.*, ante, 25.bimarginata *Say.*subplicata *Lec.*, ante, 25.punctipennis *Lec.*, ante, 25.pieta *Say.*helvines *Illiger.*var. *A. nana* Say.*Chrysomela helvines* Linn.erythropus *Lec.**Crepidodera erythropus* Mels.**Glyptina** Lec.spuria *Lec.*, ante, 26.lissotorques *Lec.*, ante, 27.**Longitarsus** Latr.nigripalpis *Lec.*, ante, 26.rubidus *Lec.*, ante, 26.

Chætoecma Steph.denticulata *Lec.**Haltica denticulata* Illiger.subviridis *Lec.*, ante, 27.**Cerotoma** (‡Chevr.).caminea *Fabr.* (Galleruca.)**Diabrotica** (‡Chevr.).thoracica *Mels.* (Calomierus).longicornis *Say* (Galleruca).trineicta *Say* (Galleruca).? atriventris *Say* (Galleruca), (fide Say).? atripennis *Say* (Galleruca), (fide Say).**Adimonia** Laich.externa *Lec.**Galleruca externa* Say.**Galleruca** Fabr.dorsata *Say* (fide Say).circumdاتا *Say* (fide Say).and five other species.¹**Hispa** Fabr.? collaris *Say* (fide Say).**Anoplitis** Kirby.scapularis *Lec.**Hispa scapularis* Oliv.*Hispa lateralis* Say.rosea *Lec.**Hispa rosea* Weber.*Hispa pallida* Say.*Hispa Philenon* Newman.**Microhophala** Chevr.lætula *Lec.*, ante, 27.cyanea *Lec.**Hispa cyanea* Say.? *Hispa Hecate* Newman.**Chelymorpha** Boheman.Argus *Boh.**Cassida Argus* Herbst.*Cassida cribraria* †Oliv.*Imitidium 17-punctatum* Say.**Cassida** Herbst.? unipunctata *Say.*? ellipsis *Lec.*, ante, 28.6-punctata *Fabr.**C. bistrispustulata* Herbst.nigripes *Oliv.*atripes *Lec.*, ante, 28.**Coptocycla** Boh.pallida *Lec.**Cassida pallida* Herbst.? *Coptocycla aurisplendens* Mann.
cruciata *Lec.**Cassida cruciata* DeG.*Cassida guttata* Oliv.*Cassida signifer* Herbst.*Coptocycla guttata* Boh.purpurata *Boh.***COCCINELLIDÆ.****Anisosticta** Chevr.vittigera *Lec.**Hippodamia vittigera* Mann.*Namia vittigera* Muls.episcopalis *Lec.**Coccinella episcopalis* Kirby.*Namia episcopalis* Muls.**Hippodamia** Chevr.glacialis *Muls.**Coccinella glacialis* Fabr.*Coccinella abbreviata* Fabr.*Coccinella remota* Weber.13-punctata *Muls.**Coccinella 13-punctata* Linn.*Coccinella tibialis* Say.Lecontii *Muls.*convergens *Guér.**Coccinella modesta* Mels.parenthesis *Lec.**Coccinella parenthesis* Say.*Coccinella tridens* Kirby.*Adonia parenthesis* Muls.**Coccinella** Linn.transversoguttata *Fald.**C. 5-notata* Kirby.monticola *Muls.*novemnotata *Herbst.*abdominalis *Say.*humeralis *Say* (fide Say).¹**Brachiacantha** Chevr.albifrons *Lec.**Coccinella albifrons* Say.tau *Lec.*, ante, 28.10-pustulata *Lec.**Hyperaspis 10-pustulata* Mels.¹ These supposed species belong to the group of pubescent trivittate ones, and need more thorough investigation than can at present be made.

Hyperaspis Chevz.

- vittigera *Lec.* Proc. Acad. Nat. Sc. VI, 133.
 quadrivittata *Lec.* *ibid.* VI, 133.
 elegans *Muls.*
Coccinella undulata||*Say.*
 pratensis *Lec.* Proc. Acad. Nat. Sc. VI, 134.

Eneis *Muls.*

- pusilla *Lec.* Proc. Acad. Nat. Sc. VI, 135.

Scymnus *Kug.*

- chatchas *Muls.*
S. collaris *Mels.*
 caudalis *Lec.*
 ? *S. creperus* *Muls.*

Sacium *Lec.*

- lunatum *Lec.* Proc. Acad. Nat. Sc. VI, 144.

II. OF EASTERN NEW MEXICO.

CICINDELIDÆ.**Amblychila** *Say.*

- cylindriciformis *Say.**
A. Piccolomini *Reiche.*

Megacephala *Latr.*

- virginica *Dej.**
Cicindela virginica *Linn.*
 carolina *Dej.*
Cicindela carolina *Linn.*

Cicindela *Linn.*

- pulchra *Say.**
 purpurea *Oliv.**
 var. *C. Audubonii* *Lec.*
 obsoleta *Say.**
 cinetipennis *Lec.**
 cuprascens *Lec.**
 guttifera *Lec.*
 punctulata *Fabr.**
 var. *C. micans* *Fabr.*
 eumatis *Lec.**
 sedecimpunctata *Klug.*

CARABIDÆ.**Micrixys** *Lec.*

- distinctus *Lec.*
Panogæus distinctus *Hald.*
Eugnathus||*distinctus* *Lec.*

Helluomorpha *Lap.*

- laticornis *Lap.**
Helluo laticornis *Dej.*

Lachnophorus *Dej.*

- elegantulus *Mann.**
Tachypus mediosignatus *Ménétr.*

Apristus *Chaud.*

- subsulcatus *Lec.*
Dromius subsulcatus *Dej.*

Glycia *Chaud.*

- viridicollis *Lec.**

Pristodactyla *Dej.*

- dubia *Lec.* Proc. Acad. Nat. Sc. VII, 38.

Platynus *Bon.* (emend. *Brullé.*)

- subeordatus *Lec.**
Agonum erythropum||*Kirby.*
 placidus *Lec.**
Feronia placida *Say.*
Agonum morosum *Dej.*

Evarthrus *Lec.*

- substriatus *Lec.**
Feronia (Molops) substriata *Lec.*

Pœcilus *Bon.*

- scitulus *Lec.**

Amara *Bon.*

- polita *Lec.**
 fareta *Lec.* Proc. Acad. Nat. Sc. VII, 353.
 musculus *Say.**
Acrodon contempta *Lec.*
 harpalina *Lec.* Proc. Acad. Nat. Sc. VII, 355.

Nothopus *Lec.*

- zabroides *Lec.**
Euryderus||*zabroides* *Lec.*
 ? *Amara grossa* *Say.*

Cratognathus *Dej.*

- setosus *Lec.**
Pisosoma setosum *Lec.*
 cordatus *Lec.* Trans. Am. Phil. Soc. X, 381.

Anisodactylus Dej.

rusticus Dej.*

Harpalus rusticus Say.*Anisodactylus pinguis* Lec.*Anisodactylus crassus* Lec.*Anisodactylus gravidus* Lec.*Anisodactylus tristis* Dej.

chalecus Lec., ante, 2.

Harpalus Latr.

impotens Lec. Journ. Ac. Nat. Sc. 2d ser. IV, 14.

amputatus Say.*

H. Stephensii Kirby.

rotundicollis Kirby.*

retractus Lec. Journ. Ac. Nat. Sc. 2d ser. IV, 29.

H. impiger Lec. Proc. Ac. Nat. Sc. VII, 79.

fallax Lec., ante, 2.

desertus Lec., ante, 3.

oblitus Lec., ante, 2.

Stenolophus Dej.

ochropezus Dej.*

Feronia ochropeza Say.

dissimilis Dej.*

Dicælus Bon.

splendidus Say.*

Chlænium Bon.

pennsylvanicus Say.*

C. vicinus Dej.*C. pubescens* Harris.

laticollis Say.*

C. diffinis Chaud.

sericeus Say.*

Carabus sericeus Forster.*Chlænium perviridis* Lec.**Pasimachus** Bon.

validus Lec.* Journ. Ac. Nat. Sc. 2d ser. IV, 14.

P. punctulatus Lec.

elongatus Lec.*

Scarites Fabr.

subterraneus Fabr.* (eum var.)

Clivina Latr.

bipustulata Dej.*

Scarites bipustulatus Fabr.*Scarites quadripustulatus* Beauv.*Clivina quadripustulata* Say.**Aspidoglossa** Putz.

subangulata Lec.*

Clivina crenata Dej.*Dyschirius subangulatus* Chaud.*Dyschirius humeralis* Chaud.*Aspidoglossa fraternata* Putz.*Aspidoglossa vicina* Putz.*Clivina bipustulata* Say.**Bembidium** Latr.

coxendix Say.*

Odontium coxendix Lec.

dorsale Say.*

pictum Lec.*

Calosoma Fabr.

scrutator Fabr.*

externum Lec.*

Carabus externus Say.*Calosoma longipenne* Dej.**DYTISCIDÆ.****Hydroporus** Clairv.

striatellus Lec.

vilis Lec.

Anisomera Brullé.

cordata Lec.

Colymbetes Clairv.

binotatus Harris.*

C. maculicollis Aubé.**Cybister** Curtis.

fimbriolatus White,* Brit. Mus. Cat. 5.

Dytiscus fimbriolatus Say.*Cybister dissimilis* Aubé.

ellipticus Lec.

HYDROPHILIDÆ.**Hydrophilus** Geoffr.

triangularis Say.*

H. lugubris Motsch.*Stelthoxus subsulcatus* Lec. Pr. Ac. VII, 221.

lateralis Fabr., Lec.* Pr. Ac. Nat. Sc. VII, 367.

ellipticus Lec. ibid. VII, 368.

STAPHYLINIDÆ.**Staphylinus** Linn.

villosus Grav.*

Philonthus Leach.

two species.

SILPHIDÆ.**Necrophorus** Fabr.mediatus *Fabr.****Silpha** Linn.peltata *Lec.***Scarabæus peltatus* Catesby.*Silpha americana* Linn.lapponica *Herbst.***S. caudata* Say.*S. tuberculata* Germ.*S. californica* Mann.*Oiceoptoma granigerum* Chevrr.truncata *Say.**ramosa *Say.****NITIDULIDÆ.****Carpophilus** Leach.pallipennis *Lec.***C. floralis* Er. Germ. Zeitschr. IV, 261.*Cercus pallipennis* Say.**Nitidula** Fabr.ziczæ *Say.****CUCUJIDÆ.****Læmophilæus** Er.biguttatus *Mels.***Cucujus biguttatus* Say.*Læmophilæus bisignatus* Guér.**Nausibius** Redt.dentatus *Redt.** Fauna Austr. 999.*Lycylus dentatus* Fabr.*Silvanus dentatus* Say.**DERMESTIDÆ.****Dermestes** Linn.marmoratus *Say.**fasciatus *Lec.* Proc. Acad. Nat. Sc. VII, 107.**Attagenus** Latr.spurens *Lec.* Proc. Acad. Nat. Sc. VII, 109.*?A. cylindricornis* Say.dichrous *Lec.* Proc. Acad. Nat. Sc. VII, 110.**Trogoderma** Latr.pusillum *Lec.* Proc. Acad. Nat. Sc. VII, 111.**PARNIDÆ.****Helichus** Er.foveatus *Lec.* Proc. Acad. Nat. Sc. VI, 43.aqualis *Lec.* Proc. Acad. Nat. Sc. VII, 81.**SCARABÆIDÆ.****Strategus** Hope.Julianus *Burm.,* Lamell. III, 133.**Aphonus** Lec.tridentatus *Lec.** Proc. Acad. Nat. Sc. VIII, 22.*Scarabæus tridentatus* Say.**Ligyryus** Burm.gibbosus *Lec.** Proc. Acad. Nat. Sc. VIII, 20.*Scarabæus gibbosus* DeGeer.*Podalgus variolosus* Burm.**Phileurus** Latr.valgus *Dej.** Burm., Lamell. III, 160.*Geotrupes valgus* Fabr.*Phileurus castaneus* Hald.**Cremastochilus** Knoch.saucius *Lec.** Journ. Ac. Nat. Sc. 2d ser. IV, 16.**Euryomia** Burm. (emend. Lac.)melancholica *Lac.** Gen. Col. III, 528.*Cetonia melancholica* Gory.Kernii *Lac.* *ibid.**Euphoria cernii* Hald.Clarkii *Lac.* *ibid.**Euphoria Clarkii* Lec. Proc. Acad. Nat. Sc. VI, 414.**Allorhina** Burm. (emend. Lac.)mutabilis? *Lac.* Gen. Col. III, 497.*Cotinis mutabilis* Burm., Lamell. I, 255.*Gymnetis mutabilis* Gory.**Anomala** Samouelle.varians *Burm.***Melolontha varians* Fabr.**Strigoderma** Burm.arboricola *Burm.***Melolontha arboricola* Fabr.**Polyphylla** Harris.decemlineata *Lec.***Melolontha 10-lineata* Say.

Thyce Lec.squamicollis *Lec.* J. Ac. Nat. Sc. 2d ser. III, 232.**Laehnosterna** Hope.lanceolata *Lec.***Melolontha lanceolata* Say.*Tostegoptera lanceolata* Blanchard.**Dichelonycha** Kirby.sulcata *Lec.* Journ. Acad. Nat. Sc. 2d ser. III, 231.**Hoplia** Illiger.laticollis *Lec.* Journ. Ac. Nat. Sc. 2d ser. III, 234.**Trox** Fabr.alternans *Lec.** Proc. Acad. Nat. Sc. VII, 211.**Canthon** Illiger.praticola *Lec.*,* ante, 10.**BUPRESTIDÆ.****Psiloptera** Sol. (emend. Lac.)Woodhousei *Lec.***Dicrera Woodhousei* Lec.var. *Psiloptera valens* Lec.**Ancylochira** Esch.confluens *Lec.***Buprestis confluens* Say.alternans *Lec.* Trans. Am. Phil. Soc. XI, 207.subornata *Lec.* Trans. Am. Phil. Soc. XI, 208.**Melanophila** Esch.miranda *Lec.**Phanops mirandus* Lec. Proc. Acad. VII, 83.**Anthaxia** Esch.retifera *Lec.* Trans. Am. Phil. Soc. XI, 215.imperfecta *Lec.* Trans. Am. Phil. Soc. XI, 215.**Chrysobothris** Esch.quadrilincata *Lec.* Trans. Am. Phil. Soc. XI, 233.cuprascens *Lec.* Trans. Am. Phil. Soc. XI, 233.**Acmæodera** Esch.variegata *Lec.* Proc. Acad. Nat. Sc. VI, 67.mixta *Lec.** Trans. Am. Phil. Soc. XI, 227.**ELATERIDÆ.****Cratonychus** Er.exuberans *Lec.* Trans. Am. Phil. Soc. X, 477.**TELEPHORIDÆ.****Silis** Charp.difficilis *Lec.***Telephorus** Geoffr.fidelis *Lec.***MELYRIDÆ.****Collops** Er.bipunctatus *Er.***Malachius bipunctatus* Say.**Dasytes** Fabr.crythropus *Lec.* Proc. Acad. Nat. Sc. VI, 171.**CLERIDÆ.****Cymatodera** Gray.longicornis *Lec.***Trichodes** Herbst.ornatus *Say.****Clerus** Geoffr.mexicanus *Spin.*maestus *Klug.**C. truncatus* Lec.**Corynetes** Fabr.rufipes *Fabr.**violaceus *Fabr.****PTINIORES.****Niptus** Boield.ventriculus *Lec.*, ante, 13.**Trypopytus** Redt.punctatus *Lec.*, ante, 13.**TENEBRIONIDÆ.****Epitragus** Latr.canaliculatus *Say.****Eurymetopon** Esch.abnorme *Lec.*atrum? *Lec.*

Pactostoma Lec.*anastomosis* Lec.**Asida anastomosis* Say.**Pelecyporus** Sol.*sordidus* Lec.* Proc. Acad. Nat. Sc. VI, 46.**Euschides** Lec.*convexicollis* Lec. Proc. Acad. Nat. Sc. VII, 224.
obovata Lec.*convexa* Lec.* ante, 14.*polita* Lec.*Asida polita* Say.**Asida** Latr.*opaca* Say.*Euschides opaca* Lec.**Zopherus** Gray.*concolor* Lec.**Eleodes** Esch.*obscura* Lec.**Blaps obscura* Say.? *Blaps hispilabris* Say.*acuta* Lec.**Blaps acuta* Say.*tricostata* Lec.**Blaps tricostata* Say.*Pimelia alternata* Kirby.*gracilis* Lec. Proc. Acad. Nat. Sc. 1858, 184.*sponsa* Lec. *ibid.* 184.*caudifera* Lec. *ibid.* 184.*obsoleta* Lec.**Blaps obsoleta* Say.*debilis* Lec. Proc. Acad. Nat. Sc. 1858, 185.*extricata* Lec.**Blaps extricata* Say.*carbonaria* Lec.**Blaps carbonaria* Say.*nigrina* Lec. Proc. Acad. Nat. Sc. 1858, 186.**Embaphion** Say.*contusum* Lec. J. Acad. Nat. Sc. 2d ser. IV, 40.***Eusattus** Lec.*reticulatus* Lec.**Zophosis reticulata* Say.**Blapstinus** Dej.*pratensis* Lec.* ante, 15.**Nyctobates** Esch.*pensylvanicus* Lec.**Tenebrio pensylvanicus* Knoch.*Upis chrysops* Herbst.**Xystropus** Sol.*pinguis* Lec., ante, 16.**Cistela** Fabr.*sericea* Say.***MELOIDÆ.****Meloe** Linn.*sublævis* Lec. Proc. Acad. Nat. Sc. VII, 84.*parvulus* Hald. Proc. Acad. Nat. Sc. VI, 404.*M. parvus*||Hald.**Lytta** Fabr.*biguttata* Lec. Proc. Acad. Nat. Sc. VI, 332.*maculata* Say.**costata* Lec. Proc. Acad. Nat. Sc. VII, 84.*longicollis* Lec.* Proc. Acad. Nat. Sc. VI, 343.*Fabricii* Lec.* Proc. Acad. Nat. Sc. VI, 343.*L. cinerea*||Fabr., Harris.**Tetraonyx** Latr.*fulva* Lec. Proc. Acad. Nat. Sc. VI, 344.**Nemognatha** Illiger.*nigripennis* Lec. Proc. Acad. Nat. Sc. VI, 347.*cribraria* Lec. Proc. Acad. Nat. Sc. VI, 348.*immaculata* Say.***Zonitis** Fabr.*flavida* Lec. Proc. Acad. Nat. Sc. VI, 349*rufa* Lec. Proc. Acad. Nat. Sc. VII, 85.**CEDEMERIDÆ.****Calopus** Fabr.*angustus* Lec.**Asclera** Schmidt.*obscura* Lec. Proc. Acad. Nat. Sc. VII, 21.**Mycterus** Oliv.*concolor* Lec. Proc. Acad. Nat. Sc. VI, 235.**CURCULIONIDÆ.****Bruchus** Linn.

one species.

Ophryastes Schönh.*tuberosus* Lec. Proc. Acad. Nat. Sc. VI, 443.*vittatus* Schönh.**Liparus vittatus* Say.

Tanymeceus Germ.
lautus *Lec.* Proc. Acad. Nat. Sc. VII, 85.

Sitones Germ.
scissifrons *Say.*

Cleonus Schönh.
lutulentus *Lec.*, ante, 18.

Anthonomus Germ.
one species.

Baridius Schönh.
one species.

Cossonus Clairv.
one species.

Tomieus Latr.
pini *Harris.**
Bostrichus pini *Say.*
caligraphus *Germ.**
Bostrichus exesus *Say.*

CERAMBYCIDÆ.

Derobrachus Serv.
geminatus *Lec.* Proc. Acad. Nat. Sc. VI, 233.

Prionus Geoffr.
curvatus *Lec.*, ante, 19.
palparis *Say.**
fissicornis *Hald.**

Criocephalus Muls.
asperatus *Lec.*, ante, 19.

Elaphidion Serv.
villosum *Hald.**
Stenocorus villosus *Fabr.*
Stenocorus putator *Peck.*
debile *Lec.* Proc. Acad. Nat. Sc. VI, 442.

Eriphus Serv.
? ignicollis *Say* (*Callidium*).
Callidium sanguinicolle *Germ.*
? rutilans *Lec.* (*Arhopalus*).

Arhopalus Serv. (emend. *Lec.*)
pietns *Lec.**
Cerambyx pictus *Drury.*
Leptura Robinie *Forster.*
Clytus flexuosus *Fabr.*

Tylosis *Lec.*
maculatus *Lec.*

Stenaspis Dupont.
solitaria *Lec.**
Cerambyx solitarius *Say.*
Smileceras solitarium *Lec.*

Rhagium *Fabr.*
lineatum *Schönh.**
Stenocorus lineatus *Oliv.*

Leptura *Linn.*
auripilis *Lec.*

Monilema *Say.*
appressum *Lec.*

Monohammus *Latr.*
clamator *Lec.*

Stenostola *Muls.*
perigrata *Lec.**
Saperda perigrata *Say.*

Tetraopes *Dalm.*
canescens *Lec.*

Saperda *Fabr.*
concolor *Lec.*

Ædilis *Serv.*
spectabilis *Lec.*, ante, 22.

CHRYSOMELINÆ.

Lema *Fabr.*
trivirgata *Lec.*, ante, 22.

Urodera *Lac.*
crucifera *Lac.* Mon. Chrysom. II, 454.

Cryptocephalus *Geoffr.*
confluens *Say.**

Colaspis *Fabr.*
one species.

Myochrous *Chevr.*
denticollis *Lec.**
Eumolpus denticollis *Say.*

Chrysomela *Linn.*
auripennis *Say.** (cum var. *caerulea*).
interrupta *Fabr.**
formosa *Say.**

Cedionychis Latr.

lugens Lec., ante, 24.

Haltica Fabr.

alternata Illiger.*

H. quinquevittata Say.

semicarbonata Lec., ante, 25.

ambiens Lec., ante, 25.

obliterata Lec., ante, 26.

punetipennis Lec.,* ante, 25.

torquata Lec., ante, 26.

biteniata Lec., ante, 26.

Luperus Geoffr.

rufipes Lec., ante, 27.

Galleruca Geoffr.

one species.

Microrhopala Chev.

cyanea Lec.*

Hispa cyanea Say.

? *Hispa Hecate* Newman.

Cassida Herbst.

sexpunctata Fabr.*

C. bistrimaculata Herbst.

EROTYLIDÆ.

Erotylus Fabr.

Boisduvalii Chev.

COCCINELLIDÆ.

Anisosticta Chev.

vittigera Lec.*

Hippodamia vittigera Mann.

Hippodamia Chev.

Lecontii Muls.*

convergens Guérin.*

H. modesta Mels.

Coccinella Linn.

monticola Mels.*

Psyllobora Chev.

vigintimaculata Say.*


Brachiacantha Chev.

albifrons Lec.*

Coccinella albifrons Say.


Epilachna Chev.

corrupta Muls. Cocc. 815.



Bayer, A. & L. 1881

- | | | | |
|------------------------|-----------------------|----|----|
| 1 DICAELUS LAEVIPENNIS | 5 LEPTOCYBUS | 9 | 13 |
| 2 CALOSOMA LUXATUM. | 6 POLYPHYLLID | 10 | 14 |
| 3 SILPHA TRUNCATA | 7 EUSAYTUS PETIOLATUS | 11 | 15 |
| 4 OMORGUS SCUTELLARIS | 8 EMBALMIDAE PRESSUM | 12 | 16 |



Boisduval

1. AMBLYCHILA CYLINDRIFORMIS. 5. THYCE SQUAMIFRONS. 9. CYSTRODEMUS VITATUS. 14. PSEPHENUS VITATUS.
 2. MICHYXIS DISTINCTUS. 6. PSILOPTERA WOODROUSEI. 10. " WISLIZENI. 15. TYRUS MASONIUS.
 3. ANSUMERA CORDATA. 7. MELANOPHILA MIRANDA. 11. PACTOSTOMA ANASTOMOSIS. 16. PACTOSTOMA SPECIOSUM.
 4. PSEPHENUS VALGUS. 8. PROTYLUS BOISDUVALII. 12. DEROBRACHUS GEMINATUS. 17. MONTICOMA APPRESSUM.
 13. PRICNUS EMARGINATUS.



THIS BOOK IS DUE ON THE LAST DATE
STAMPED BELOW

AN INITIAL FINE OF 25 CENTS
WILL BE ASSESSED FOR FAILURE TO RETURN
THIS BOOK ON THE DATE DUE. THE PENALTY
WILL INCREASE TO 50 CENTS ON THE FOURTH
DAY AND TO \$1.00 ON THE SEVENTH DAY
OVERDUE.

BIOLOGY LIBRARY

SEP 9 1941

DEC 11 1941

NOV 2 1948

NOV 22 1948

NOV 8 1950

MAY 27 1952

MAY 15 1952

MAR 16 1984

LD 21-10m-7.39 (402s)

Coleoptera of Kansas and
eastern New Mexico.

K3L4

BIOLOGY
LIBRARY
6

Bio. Lib.

SEP 9 1941 *Wick* SEP 12 1941
DEC 11 1941 *Wick* NOV 27 1941

U.C. BERKELEY LIBRARIES



039321301

299243 *QL584*

K3L4

Leconte

BIOLOGY
LIBRARY
6

UNIVERSITY OF CALIFORNIA LIBRARY

